



Cochrane
Library

Cochrane Database of Systematic Reviews

Hyperbaric oxygen therapy for acute coronary syndrome (Review)

Bennett MH, Lehm JP, Jepson N

Bennett MH, Lehm JP, Jepson N.

Hyperbaric oxygen therapy for acute coronary syndrome.

Cochrane Database of Systematic Reviews 2015, Issue 7. Art. No.: CD004818.

DOI: 10.1002/14651858.CD004818.pub4.

www.cochranelibrary.com

TABLE OF CONTENTS

HEADER	1
ABSTRACT	1
PLAIN LANGUAGE SUMMARY	2
SUMMARY OF FINDINGS FOR THE MAIN COMPARISON	3
BACKGROUND	4
OBJECTIVES	5
METHODS	5
RESULTS	7
Figure 1.	8
Figure 2.	11
ADDITIONAL SUMMARY OF FINDINGS	13
DISCUSSION	16
AUTHORS' CONCLUSIONS	17
ACKNOWLEDGEMENTS	17
REFERENCES	18
CHARACTERISTICS OF STUDIES	20
DATA AND ANALYSES	28
Analysis 1.1. Comparison 1 Death, Outcome 1 Death at any time.	30
Analysis 1.2. Comparison 1 Death, Outcome 2 Death - best case scenario.	31
Analysis 1.3. Comparison 1 Death, Outcome 3 Death - worst case scenario.	32
Analysis 2.1. Comparison 2 Major Adverse Cardiac Events, Outcome 1 Major Adverse Cardiac Events.	33
Analysis 2.2. Comparison 2 Major Adverse Cardiac Events, Outcome 2 MACE - Best case scenario.	33
Analysis 2.3. Comparison 2 Major Adverse Cardiac Events, Outcome 3 MACE - worst case scenario.	34
Analysis 2.4. Comparison 2 Major Adverse Cardiac Events, Outcome 4 Recurrent acute myocardial infarction (AMI).	34
Analysis 2.5. Comparison 2 Major Adverse Cardiac Events, Outcome 5 Recurrent AMI - best case scenario.	35
Analysis 2.6. Comparison 2 Major Adverse Cardiac Events, Outcome 6 Recurrent AMI - Worst case scenario.	35
Analysis 3.1. Comparison 3 Significant dysrhythmias (complete heart block, ventricular fibrillation, asystole), Outcome 1 Overall (CHB, VF and asystole combined).	36
Analysis 3.2. Comparison 3 Significant dysrhythmias (complete heart block, ventricular fibrillation, asystole), Outcome 2 Significant dysrhythmias (complete heart block, ventricular fibrillation or asystole).	37
Analysis 3.3. Comparison 3 Significant dysrhythmias (complete heart block, ventricular fibrillation, asystole), Outcome 3 Overall best case.	38
Analysis 3.4. Comparison 3 Significant dysrhythmias (complete heart block, ventricular fibrillation, asystole), Outcome 4 Overall worst case.	38
Analysis 4.1. Comparison 4 Time to pain relief, Outcome 1 Time to relief of pain.	39
Analysis 5.1. Comparison 5 Magnitude of cardiac enzyme changes, Outcome 1 12 hour Plasma Creatine Phosphokinase.	39
Analysis 5.2. Comparison 5 Magnitude of cardiac enzyme changes, Outcome 2 24 hour Plasma Creatine Phosphate.	40
Analysis 5.3. Comparison 5 Magnitude of cardiac enzyme changes, Outcome 3 Maximum Plasma Creatine Phosphate.	40
Analysis 6.1. Comparison 6 Improvement in left ventricular function, Outcome 1 Improved contraction in at least one segment (post-HBOT echo).	41
Analysis 6.2. Comparison 6 Improvement in left ventricular function, Outcome 2 Left Ventricular Ejection Fraction- % (final estimate).	41
Analysis 7.1. Comparison 7 Length of Stay, Outcome 1 Overall length of stay (days).	42
Analysis 8.1. Comparison 8 Adverse events of therapy, Outcome 1 Total adverse events.	42
APPENDICES	43
WHAT'S NEW	56
HISTORY	57
CONTRIBUTIONS OF AUTHORS	57
DECLARATIONS OF INTEREST	57
SOURCES OF SUPPORT	58
INDEX TERMS	58

[Intervention Review]

Hyperbaric oxygen therapy for acute coronary syndrome

Michael H Bennett¹, Jan P Lehm², Nigel Jepson³

¹Department of Anaesthesia, Prince of Wales Clinical School, University of NSW, Sydney, Australia. ²Department of Diving and Hyperbaric Medicine, Prince of Wales Hospital, Randwick, Australia. ³Prince of Wales Hospital, Randwick, Australia

Contact address: Michael H Bennett, Department of Anaesthesia, Prince of Wales Clinical School, University of NSW, Sydney, NSW, Australia. m.bennett@unsw.edu.au. s9400356@unsw.edu.au.

Editorial group: Cochrane Heart Group.

Publication status and date: New search for studies and content updated (no change to conclusions), published in Issue 7, 2015.

Review content assessed as up-to-date: 17 September 2014.

Citation: Bennett MH, Lehm JP, Jepson N. Hyperbaric oxygen therapy for acute coronary syndrome. *Cochrane Database of Systematic Reviews* 2015, Issue 7. Art. No.: CD004818. DOI: 10.1002/14651858.CD004818.pub4.

Copyright © 2015 The Cochrane Collaboration. Published by John Wiley & Sons, Ltd.

ABSTRACT

Background

Acute coronary syndrome (ACS), includes acute myocardial infarction and unstable angina, is common and may prove fatal. Hyperbaric oxygen therapy (HBOT) will improve oxygen supply to the threatened heart and may reduce the volume of heart muscle that perishes. The addition of HBOT to standard treatment may reduce death rate and other major adverse outcomes.

This is an update of a review previously published in May 2004 and June 2010.

Objectives

The aim of this review was to assess the evidence for the effects of adjunctive HBOT in the treatment of ACS. We compared treatment regimens including adjunctive HBOT against similar regimens excluding HBOT. Where regimens differed significantly between studies this is clearly stated and the implications discussed. All comparisons were made using an intention to treat analysis where this was possible. Efficacy was estimated from randomised trial comparisons but no attempt was made to evaluate the likely effectiveness that might be achieved in routine clinical practice. Specifically, we addressed:

Does the adjunctive administration of HBOT to people with acute coronary syndrome (unstable angina or infarction) result in a reduction in the risk of death?

Does the adjunctive administration of HBOT to people with acute coronary syndrome result in a reduction in the risk of major adverse cardiac events (MACE), that is: cardiac death, myocardial infarction, and target vessel revascularization by operative or percutaneous intervention?

Is the administration of HBOT safe in both the short and long term?

Search methods

We updated the search of the following sources in September 2014, but found no additional relevant citations since the previous search in June 2010 (CENTRAL), MEDLINE, EMBASE, CINAHL and DORCTHIM. Relevant journals were handsearched and researchers in the field contacted. We applied no language restrictions.

Selection criteria

Randomised studies comparing the effect on ACS of regimens that include HBOT with those that exclude HBOT.

Hyperbaric oxygen therapy for acute coronary syndrome (Review)

Copyright © 2015 The Cochrane Collaboration. Published by John Wiley & Sons, Ltd.

Data collection and analysis

Three authors independently evaluated the quality of trials using the guidelines of the Cochrane Handbook and extracted data from included trials. Binary outcomes were analysed using risk ratios (RR) and continuous outcomes using the mean difference (MD) and both are presented with 95% confidence intervals. We assessed the quality of the evidence using the GRADE approach.

Main results

No new trials were located in our most recent search in September 2014. Six trials with 665 participants contributed to this review. These trials were small and subject to potential bias. Only two reported randomisation procedures in detail and in only one trial was allocation concealed. While only modest numbers of participants were lost to follow-up, in general there is little information on the longer-term outcome for participants. Patients with acute coronary syndrome allocated to HBOT were associated with a reduction in the risk of death by around 42% (RR: 0.58, (95% CI 0.36 to 0.92), 5 trials, 614 participants; low quality evidence).

In general, HBOT was well-tolerated. No patients were reported as suffering neurological oxygen toxicity and only a single patient was reported to have significant barotrauma to the tympanic membrane. One trial suggested a significant incidence of claustrophobia in single occupancy chambers of 15% (RR of claustrophobia with HBOT 31.6, 95% CI 1.92 to 521).

Authors' conclusions

For people with ACS, there is some evidence from small trials to suggest that HBOT is associated with a reduction in the risk of death, the volume of damaged muscle, the risk of MACE and time to relief from ischaemic pain. In view of the modest number of patients, methodological shortcomings and poor reporting, this result should be interpreted cautiously, and an appropriately powered trial of high methodological rigour is justified to define those patients (if any) who can be expected to derive most benefit from HBOT. The routine application of HBOT to these patients cannot be justified from this review.

PLAIN LANGUAGE SUMMARY

Does hyperbaric oxygen therapy improve outcome after heart attack?

Background

Acute heart attacks and severe angina (heart pain) are usually due to blockages in the arteries supplying the heart (coronary arteries). These problems are collectively referred to as 'acute coronary syndrome' (ACS). ACS is very common and may lead to severe complications including death. Hyperbaric oxygen therapy (HBOT) involves people breathing pure oxygen at high pressures in a specially designed chamber. It is sometimes used as a treatment to increase the supply of oxygen to the damaged heart in an attempt to reduce the area of the heart that is at risk of dying.

We searched the medical literature for any studies that reported the outcome of patients with ACS when treated with HBOT.

Studies found

We first searched the literature in 2004 and most recently in September 2014, finding 6 studies in total. All studies included patients with heart attack and some also included patients with severe angina. The dose of hyperbaric oxygen was similar in most studies.

Key results

Overall, we found some evidence that people with ACS are less likely to die or to have major adverse events, and to have more rapid relief from their pain if they receive hyperbaric oxygen therapy as part of their treatment. However, our conclusions are based on relatively small randomised trials. Our confidence in these findings is further reduced because in most of these studies both the patients and researchers were aware of who was receiving HBOT and it is possible a 'placebo effect' has biased the result in favour of HBOT. HBOT was generally well-tolerated. Some patients complained of claustrophobia when treated in small (single person) chambers and there was no evidence of important toxicity from oxygen breathing in any subject. One individual suffered damage to the eardrum from pressurisation.

Conclusions

While HBOT may reduce the risk of dying, time to pain relief and the chance of adverse heart events in people with heart attack and unstable angina, more work is needed to be sure that HBOT should be recommended.

SUMMARY OF FINDINGS FOR THE MAIN COMPARISON *[Explanation]*

hyperbaric oxygen therapy for acute coronary syndrome						
Patient or population: patients with acute coronary syndrome Settings: Acute care hospital Intervention: hyperbaric oxygen therapy						
Outcomes	Illustrative comparative risks* (95% CI)		Relative effect (95% CI)	No of Participants (studies)	Quality of the evidence (GRADE)	Comments
	Assumed risk	Corresponding risk				
	Control	hyperbaric oxygen therapy				
Death at any time	Study population		RR 0.58 (0.36 to 0.92)	614 (5 studies)	⊕⊕○○ low ^{1,2}	
	116 per 1000	67 per 1000 (42 to 107)				
	Medium risk population					
	102 per 1000	59 per 1000 (37 to 94)				
<p>*The basis for the assumed risk (e.g. the median control group risk across studies) is provided in footnotes. The corresponding risk (and its 95% confidence interval) is based on the assumed risk in the comparison group and the relative effect of the intervention (and its 95% CI).</p> <p>CI: Confidence interval; RR: Risk ratio;</p>						
GRADE Working Group grades of evidence						
High quality: Further research is very unlikely to change our confidence in the estimate of effect.						
Moderate quality: Further research is likely to have an important impact on our confidence in the estimate of effect and may change the estimate.						
Low quality: Further research is very likely to have an important impact on our confidence in the estimate of effect and is likely to change the estimate.						
Very low quality: We are very uncertain about the estimate.						

¹ Critical outcome

² Small sample with low numbers of events

BACKGROUND

Description of the condition

Cardiovascular disease (CVD) is the leading cause of death in the world, accounting for 30% of all deaths, of which 42% are due to coronary heart disease (CHD) (WHO 2013). In the United Kingdom, CVD is the most common cause of premature death, causing just over 159,000 deaths, approximately 25% of all deaths episodes in 2011 (BHF 2012). Of these deaths, nearly 74,000 were due to CHD at a community cost of around GBP 1.8 billion in 2009 (Nichols 2012). Since myocardial infarction (the presence of two out of three of: chest pain, ECG changes and cardiac enzyme rise) is not always diagnosable during an acute event, unstable or persisting ischaemic heart pain (angina) with or without infarction are together described as acute coronary syndrome (ACS). The main underlying problems in coronary heart disease is atherosclerosis, a degenerative process characterised by the formation of plaques comprised of platelets, cells, matrix fibres, lipids, and tissue debris in the vessel lumen. While such plaques are often complicated by ulceration of the vessel wall with obstruction to blood flow, such ulceration is not necessary for plaques to be problematic (Naghavi 2003). An unstable plaque (coronary atheroma vulnerable to rupture and fissure, and associated with thrombus formation) can lead to an acute coronary syndrome without the artery being totally occluded and infarction may follow (Heistad 2003). A significant proportion of patients admitted with acute myocardial infarction (AMI) will suffer major morbidity or mortality, even when thrombolysis or angioplasty is used to relieve the obstruction, although there is some evidence that the rate of in-hospital events and six-month readmissions are falling with increasing adherence to evidence-based guidelines (Kalla 2006; Aliprandi-Costa 2011).

Description of the intervention

Hyperbaric oxygen therapy (HBOT) is an adjunctive therapy that has been proposed to improve outcome following ACS. HBOT is the therapeutic administration of 100% oxygen at environmental pressures greater than one atmosphere absolute (ATA), and involves placing the patient in an airtight vessel, increasing the pressure within that vessel, and administering 100% oxygen for respiration. In this way, it is possible to deliver a greatly increased partial pressure of oxygen to the tissues. At 2 ATA, for example, patients with reasonable cardiopulmonary function will have an arterial oxygen tension of over 1000 mmHg, and a muscle oxygen tension around 221 mmHg (Sheffield 1988; Wells 1977). In comparison, muscle oxygen tension on air at 1ATA is about 29 mmHg and 59 mmHg breathing 100% oxygen at 1ATA. Indeed, at 3ATA on 100% oxygen, there are more than 6 mls of oxygen dissolved in every 100 mls of plasma, enough to sustain basal metabolic requirements without any oxygen transport by haemoglobin (Boerema

1960; Hammarlund 1999; Leach 1998). However 3ATA of 100% oxygen becomes rapidly toxic in the brain, manifested in epileptiform grand mal seizures. Therefore in practice, treatments involve pressurisation to between 1.5 and 2.8 ATA for periods between 60 and 120 minutes once or more daily.

HBOT for ACS was first reported in a canine experimental model in 1958 (Smith 1958) and in a human subject in 1964 (Moon 1964). Several uncontrolled human studies have been published since that time, generally with indications of benefit measured as a reduction in mortality or improvements in haemodynamic or metabolic parameters (Ashfield 1969; Kline 1970). As far as the authors are aware, however, HBOT is not in routine use anywhere for patients presenting with ACS.

How the intervention might work

The administration of HBOT is based on the argument that the myocardium is hypoxic, and that HBOT can reverse that hypoxia in areas that are marginally perfused. This effect is achieved by greatly increasing the diffusion gradient down which oxygen moves from the blood to the myocyte. Improved oxygen availability may also improve outcome through the effects of oxygen as a modulator of tissue repair. Oxygen has been shown to increase the expression of antioxidant enzymes in both tissues and plasma through an increase in glutathione levels (Harabin 1990; Speit 2000), to reduce the degree of lipid peroxidation (Thom 1991) and to prevent the activation of neutrophils in response to endothelial damage, thus modifying ischaemia-reperfusion injury (Jones 2010; Tjarnstrom 1999). HBOT also mobilises stem cells from the bone marrow in a dose-dependent manner and may be important in neovascularisation of healing tissue (Heyboer 2014). The induction of protective mechanisms via a degree of oxidative stress is probably the common factor for many of these beneficial effects (Thom 2009). However despite more than 40 years of interest in the delivery of HBOT relatively little clinical evidence exists for the assertion that such an intervention improves outcome.

Why it is important to do this review

To our knowledge, there are no other systematic reviews of the clinical use of HBOT for ACS, and it is important to clearly assess both the risks and benefits of this treatment. While HBOT may produce benefit for the myocardium via the mechanisms outlined above, HBOT is associated with some risk of adverse effects including damage to the ears, sinuses and lungs from the effects of pressure (Shupak 2008), temporary worsening of short-sightedness (Khan 2003), claustrophobia and oxygen poisoning (Butler 2008; Leach 1998). Reported rates vary widely but, for example, about 20% of patients will experience some degree of middle ear barotrauma, and 60% to 70% a measurable worsening of short-sightedness (Khan 2003; Shupak 2008). Oxygen poisoning may

occur acutely in the form of grand mal seizures while exposed to hyperbaric oxygen (acute neurological toxicity), or develop over the course of treatment, resulting in a reversible reduction in vital capacity and other respiratory indices (pulmonary oxygen toxicity) (Clark 2008). In addition, the occurrence of significant post-infarction events such as malignant arrhythmia, might be associated with a worse prognosis if they arise while the patient is confined in a hyperbaric chamber. Although serious adverse events are rare, HBOT cannot be regarded as an entirely benign intervention (Leach 1998). For a number of reasons, therefore, the administration of HBOT for acute coronary syndrome patients remains controversial.

OBJECTIVES

The aim of this review was to assess the evidence for the effects of adjunctive HBOT in the treatment of ACS. We compared treatment regimens including adjunctive HBOT against similar regimens excluding HBOT. Where regimens differed significantly between studies this is clearly stated and the implications discussed. All comparisons were made using an intention to treat analysis where this was possible. Efficacy was estimated from randomised trial comparisons but no attempt was made to evaluate the likely effectiveness that might be achieved in routine clinical practice. Specifically, we addressed:

- Does the adjunctive administration of HBOT to people with acute coronary syndrome (unstable angina or infarction) result in a reduction in the risk of death?
- Does the adjunctive administration of HBOT to people with acute coronary syndrome result in a reduction in the risk of major adverse cardiac events (MACE), that is: cardiac death, myocardial infarction, and target vessel revascularization by operative or percutaneous intervention?
- Is the administration of HBOT safe in both the short and long term?

METHODS

Criteria for considering studies for this review

Types of studies

Randomised controlled trials that compare the effect of treatment for ACS (including thrombolysis) where HBOT administration is included, with the effect of similar treatment in the absence of

HBOT. Studies were considered irrespective of the use of a sham therapy, allocation concealment or blinding status.

Types of participants

Any adult with an acute coronary syndrome, with or without S-T segment elevation.

Types of interventions

HBOT administered in a compression chamber between pressures of 1.3ATA and 3.0ATA and treatment times between 30 minutes and 120 minutes on at least one occasion, were eligible. We accepted any standard treatment regimen designed to maximise recovery, and where the same regimen is delivered in both arms of any single trial. Subgroup analysis was intended to evaluate the impact of different comparator strategies.

Types of outcome measures

Studies were eligible for inclusion if they reported any of the following outcome measures at any time.

Primary outcomes

1. Death rate at any time following presentation;
2. Rate of Major Adverse Cardiac Events (MACE), this includes death, recurrent MI, urgent revascularization (CABG or PTCA).

Secondary outcomes

1. Rate of significant cardiac events including dysrhythmia, onset of cardiac failure, haemodynamic change;
2. Time to relief of cardiac pain;
3. Size of infarct area;
4. Magnitude of cardiac enzyme changes;
5. Left ventricular function;
6. Length of stay in either a specialist cardiac unit or general hospital ward;
7. Myocardial perfusion measured by whatever means;
8. Quality of life (QOL);
9. Rate of re-admission;
10. Costs for the delivery of care;
11. Adverse events associated with HBOT including damage to the ears, sinuses and lungs from the effects of pressure, worsening of myopia, claustrophobia and oxygen poisoning. Any other adverse events reported in either arm will also be recorded.

Search methods for identification of studies

It was our intention to capture both published and unpublished studies. We applied no language restrictions.

Electronic searches

Initial searches were made in May 2004 ([Appendix 1](#)) and repeated with some modifications in strategy in June 2007 ([Appendix 2](#)), June 2010 ([Appendix 3](#)) and September 2014 ([Appendix 4](#)).

We searched: the Cochrane Central Register of Controlled Trials (CENTRAL, Issue 8, September 2014) on the *Cochrane Library*, OVID MEDLINE (1966 to September 2014), OVID EMBASE (1980 to September 2014), EBSCO CINAHL (1982 to September 2014), LILACS on Bireme (1980 to September 2014) and an additional database developed in our Hyperbaric facility, DORCTHIM (The Database of Randomised Trials in Hyperbaric Medicine, [Bennett 2002](#) - searched September 2014). The LILACS and DORCTHIM searches were by the keywords 'coronary or cardiac or heart or myocard\$' and 'hyperbaric oxygen\$'.

Searching other resources

In addition we undertook a systematic search for relevant controlled trials in specific hyperbaric literature sources up to September 2014:

1. Experts in the field and leading hyperbaric therapy centres (as identified by personal communication and searching the Internet) were contacted and asked for additional relevant data in terms of published or unpublished randomised trials;
2. Handsearching of relevant hyperbaric textbooks ([Jain 2009](#); [Kindwall 2008](#); [Mathieu 2006](#)), journals (Undersea and Hyperbaric Medicine, Hyperbaric Medicine Review, South Pacific Underwater Medicine Society (SPUMS) Journal, European Journal of Hyperbaric Medicine and Aviation, Space and Environmental Medicine Journal) and conference proceedings (Undersea and Hyperbaric Medical Society, SPUMS, European Undersea and Baromedical Society, International Congress of Hyperbaric Medicine) published since 1980;
3. Contact with authors of relevant studies to request details of unpublished or ongoing investigations.
4. We also searched for ongoing relevant trials in the registry [ClinicalTrials.gov](#) using the term 'hyperbaric oxygen'. Authors were contacted if there was any ambiguity about the published data.

Data collection and analysis

Selection of studies

One author (MB) was responsible for handsearching and identification of appropriate studies for consideration. Two authors (MB and NJ) examined the electronic search results and identified studies that were possibly relevant. All studies considered possibly relevant by at least one author were entered into a bibliographic software package (Reference Manager). All comparative clinical trials identified were retrieved in full and reviewed independently

by three authors, two with content expertise with HBOT and one with content expertise in treating ACS. In addition one of the authors (MB) has expertise in clinical epidemiology.

Data extraction and management

Using a data extraction form developed for this review, each author extracted relevant data. Any disagreements were resolved by consensus and communication with the authors of the original trials as appropriate. All data extracted reflected original allocation group where possible to allow an intention to treat analysis. Withdrawals were identified where this information was given.

Assessment of risk of bias in included studies

We followed the guidelines of the Cochrane Handbook for Systematic Reviews of Interventions for assessing the risk of bias in included studies ([Higgins 2008](#)). Two authors (MB and JL) independently assessed the quality of the studies with respect to sequence generation, allocation concealment, blinding, incomplete outcome data, selective outcome reporting and other potential threats to validity. We assessed the quality factors of each study separately and have presented them in a Risk of Bias table for each study.

Any differences of opinion were resolved by discussion and consensus.

Dealing with missing data

We employed sensitivity analyses using different approaches to impute missing data. The best-case scenario assumed that none of the originally enrolled patients missing from the primary analysis in the treatment group had the negative outcome of interest whilst all those missing from the control group did. The worst-case scenario was the reverse.

Assessment of heterogeneity

For dichotomous outcomes relative risk (RR) was used. We used a fixed-effect model where there was no evidence of significant heterogeneity between studies, and a random-effects model where inter-study heterogeneity was likely (see below under subgroups).

Data synthesis

Primary outcomes

1. Risk of death (time of outcome was determined by trial data). The RR for survival with HBOT was established using the intention to treat data of the HBOT versus the control group. Where there were withdrawals without an indication as to group allocation, we divided them in the proportions intended by randomisation. Analyses were performed with RevMan 5.0.23

software. As an estimate of the statistical significance of a difference between experimental interventions and control interventions we calculated RR, with 95% confidence intervals (CI), for survival using HBOT. A statistically significant difference between experimental intervention and control intervention was assumed if the 95% CI of the RR did not include the value 1.0. As an estimate of the clinical relevance of any difference between experimental intervention and control intervention we calculated the number-needed-to-treat (NNT) with 95% CI as appropriate;

2. Risk of suffering Major Adverse Cardiac Event (MACE - includes death, recurrent MI and urgent revascularization (CABG or PTCA)). The RR for MACE with and without HBOT was calculated using the methods described in (1) above;

Secondary outcomes

1. Risk of suffering significant cardiac events (dysrhythmia requiring intervention, haemodynamic disturbance requiring intervention or cardiac failure). The RR for cardiac event with and without HBOT was calculated using the methods described in (1) above;

2. Time to relief of cardiac pain: the weighted mean differences (WMD) in time to relief between HBOT and control groups was compared using RevMan 4.2. A statistically significant difference was defined as existing if the 95% CI did not include a zero WMD;

3. Infarct area: the WMD in infarct area was to be compared using the methods described in (4) above;

4. Rise in cardiac enzymes: the WMD between groups for the maximum enzyme level was compared as for (4) above;

5. Left ventricular function: the WMD of ejection fraction or other measure of left ventricular function was compared as for (4) above;

6. Length of Stay: the WMD in length of stay in both intensive care area/coronary care area and general hospital ward was calculated in a way analogous to that described in (4) above;

7. Myocardial perfusion: the WMD in measures designed to assess myocardial perfusion were to be compared as for (4) above;

8. Quality of life: WMD in QOL measures and/or activities of daily living were to be compared as for (4) above;

9. Risk of re-admission following treatment for acute coronary syndrome: the RR for re-admission following HBOT and comparator was to be compared using the methods described in (1) above;

10. Cost: the WMD in costs between treatment arms was to be compared as for (4) above;

11. Dichotomous data were considered for adverse events (number of patients with adverse events versus number of patients without them in both groups) in the HBOT groups of the included studies. We tabulated any recorded adverse events and pooled as appropriate.

Subgroup analysis and investigation of heterogeneity

We considered, but excluded, subgroup analysis based on:

1. Inclusion or otherwise of thrombolysis in both arms of the trial;
2. Nature of comparator treatment modalities;
3. Dose of oxygen received (pressure, time and length of treatment course);
4. Presence or absence of cardiac failure;
5. Site of infarct;
6. Infarcted subjects versus pre-infarction subjects.

Subgroup analysis by sex and age was not considered in this analysis because we did not intend to seek individual patient data. Clinical and statistical heterogeneity was explored and subgroup analyses would have been performed if appropriate. Clinically we considered differences in patient groups, the timing and nature of all therapies and other aspects of the clinical setting. Statistically, the forest plots generated were examined and the presence or absence of overlap in the confidence intervals noted (lack of overlap of confidence intervals may indicate heterogeneity). Statistical heterogeneity was to be assumed if the I^2 value exceeded 30%, and consideration would have been given to the appropriateness of pooling and meta-analysis.

Sensitivity analysis

We intended to perform sensitivity analyses for missing data and study quality where appropriate data existed.

RESULTS


Description of studies

Results of the search

Our previous searches from May 2004 to June 2010 produced a total of 158 records. After removal of duplicates, 44 records remained. On the basis of screening the titles and abstracts, we excluded 28 records, leaving 16 records that were examined in full. Of these, we included 11 reports of six trials. Our most recent search in September 2014 retrieved a further 66 records. After removal of duplicates, 38 records remained. After screening the titles and abstracts, we excluded all these records. This latest search has therefore uncovered no further trials that contribute to either the qualitative or quantitative review.

The results of all four searches are combined and summarised in Figure 1. In total we have included eleven reports of six trials (Dekleva 2004; Dotsenko 2007; Hot MI 1998; Sharifi 2004; Swift 1992; Thurston 1973).

Figure 1. Study flow diagram



Included studies

This review includes eleven reports of six trials (Dekleva 2004; Dotsenko 2007; Hot MI 1998; Sharifi 2004; Swift 1992; Thurston 1973). The secondary citations for these studies are listed in [Included studies](#). The six included trials were published between 1973 and 2007 and from a wide range of geographical locations (USA: Long Beach California Hot MI 1998 and Cleveland Ohio Sharifi 2004; Russia: Minsk Dotsenko 2007; The Balkans: Belgrade, Serbia and Montenegro Dekleva 2004; England: London Thurston 1973 and Australia: Fremantle Swift 1992). None of the trials declared any industry funding, although Thurston 1973 declared the chamber manufacturer loaned his group the hyperbaric chambers used. The authors of this review are unaware of any ongoing RCTs in the area. In total, the six trials enrolled 665 participants, 337 receiving HBOT and 328 control. The largest trial (Thurston 1973) accounts for 33% of cases. (See [Characteristics of included studies](#)).

All studies involved the administration of 100% oxygen at between 2ATA and 3ATA for between 30 and 120 minutes, however the total number of treatment sessions varied between studies. The lowest number administered was a single session (Dekleva 2004; Hot MI 1998; Swift 1992), while the highest was a maximum of 16 treatments within 48 hours (Thurston 1973).

All trials included participants with acute myocardial infarction and Sharifi 2004 also included individuals presenting with unstable angina. All trials included patients with similar age and sex distribution (see [Characteristics of included studies](#)). Only Swift 1992 described allocation concealment and blinded subjects to allocation with a sham HBOT session. The time from presentation to enrolment varied from “up to ten days” (Dotsenko 2007), “within one week” (Swift 1992) to “within 24 hours” (Thurston 1973) and “within six hours” (Hot MI 1998). The mean time to treatment with HBOT was 13 hours in Dekleva 2004, whilst Sharifi 2004 did not state any time. The primary purpose of five of these reports was the treatment of AMI with HBOT (Dotsenko 2007 was aimed at the prevention of further AMI rather than acute treatment), while for Swift 1992 it was the use of HBOT in AMI patients to identify myocardial segments capable of functional improvement, and for Sharifi 2004 the effect of HBOT on re-stenosis following percutaneous coronary interventions. Specific exclusion criteria varied between trials. All trials excluded those unfit for HBOT, but in addition Hot MI 1998 and Dekleva 2004 excluded subjects who were not suitable for thrombolysis (e.g. recent stroke), those with previous AMI and those in cardiogenic shock, while Swift 1992 excluded those with uncontrolled heart failure and/or significant ongoing angina. Thurston 1973 excluded subjects over 70 years and those presenting when there was no HBOT

chamber available. Dekleva 2004 also excluded patients over 70 or with heart failure, significant dysrhythmia or no definitive diagnosis of AMI. Sharifi 2004 excluded those who continued to show evidence of ischaemia after 30 minutes of medical treatment. Dotsenko 2007 only enrolled patients more than three days after AMI.

Comparator therapies also varied between trials. All trials employed HBOT as an adjunctive procedure to “standard” care: Hot MI 1998 and Dotsenko 2007 used thrombolysis, aspirin, heparin and intravenous nitroglycerine in suitable patients, Sharifi 2004 employed stenting and a regimen using aspirin, heparin and clopidogrel, Dekleva 2004 used thrombolysis, while Thurston 1973 used “full orthodox care” and Swift 1992 used “customary care”. The follow-up periods varied between the period immediately following HBOT (Swift 1992), to three weeks (Thurston 1973), six weeks (Dekleva 2004) eight months (Sharifi 2004) and two years (Dotsenko 2007). Hot MI 1998 reported mortality to discharge from hospital. All included studies reported at least one clinical outcome of interest. Of the outcomes identified above, these trials reported data on both primary outcomes (mortality and MACE), but only length of stay, time to pain relief, magnitude of cardiac enzyme changes, left ventricular function and adverse events from the secondary outcomes of interest.

Other outcomes reported included: angiographic re-stenosis and recurrence of angina (Sharifi 2004), left ventricular wall motion score index, diastolic filling and left ventricular end diastolic and systolic volumes (Dekleva 2004), and left ventricular ejection fraction and resolution of ST segment abnormality (Hot MI 1998).

Excluded studies

A total of five studies were excluded after review of the full report. Details are given in the table [Characteristics of excluded studies](#). Two were reports of animal experiments (Ciocatto 1965; Thomas 1990), one was a case series (Cameron 1965), one was a non-random comparative trial (Dai 1995) and one was an RCT which included patients with both stable angina and ACS, and for which the results of those with ACS could not be separately identified (Markarian 1991).

There were no ongoing studies of relevance at the registry ClinicalTrials.gov.

Risk of bias in included studies

Details of the quality assessment are given in the [Characteristics of included studies](#) table. The significance of variations in quality detailed below is unclear and given that few analyses could be pooled, study quality was not used as a basis for sensitivity analysis.

Randomisation

Randomisation procedures were described in [Hot MI 1998](#) and [Dotsenko 2007](#) (random number tables and computer-generated sequence respectively) but not in the other studies. Allocation concealment was adequately described only by [Swift 1992](#). For none of the remaining studies is there a clear indication that the investigators were unable to predict the prospective group to which a participant would be allocated.

Patient baseline characteristics

All patients required a clinical diagnosis of AMI for enrolment in these studies except [Sharifi 2004](#), who also enrolled subjects with unstable angina. In general, there were no potentially important differences in baseline characteristics between groups within each study with the exception of [Dekleva 2004](#). In that study, there were more diabetic patients in the hyperbaric group (22% versus 5%). Three studies defined entry criteria as those patients with a clinical diagnosis of AMI, [Dekleva 2004](#) within the time frame for thrombolysis, [Hot MI 1998](#) within six hours, [Thurston 1973](#) within 24 hours and [Dotsenko 2007](#) from three to seven days after the event. [Swift 1992](#) enrolled patients with AMI and abnormal left ventricular wall motion between three and seven days post-infarct. All patients in the [Sharifi 2004](#) study had presumed coronary arterial lesions where a percutaneous stent was indicated and so were a more highly selected subset of ACS patients. [Hot MI 1998](#), [Dekleva 2004](#) and [Swift 1992](#) indicated that patients who were unstable or in gross left ventricular failure were excluded.

Blinding

Only [Swift 1992](#) described the use of a sham therapy to blind participants as to treatment group allocation. This paper also described a blinded and randomised method for outcome assessment, including the measurement of concordance between multiple assessors. The echocardiographers were blinded in [Dekleva 2004](#).

Patients lost to follow-up

The percentage of patients lost to follow up in the six trials ranged from 0% to 13%. [Swift 1992](#) and [Dekleva 2004](#) reported no losses to follow-up or any violation of treatment protocol. [Hot MI 1998](#) reported 16 subjects (12%) withdrawn from analysis after allocation to groups (four became unstable, four generated incomplete data, three were enrolled after six hours in violation of inclusion criteria, two showed no cardiac enzyme rise, two received an incorrect treatment protocol and one refused to have HBOT). [Thurston 1973](#) similarly did not report data on 13 subjects (6%) who were withdrawn for misdiagnosis or being aged more than 70 years in violation of inclusion criteria. The group allocation was not indicated for any of the withdrawn patients in either of

these studies. [Sharifi 2004](#) excluded nine subjects (13%) allocated to HBOT from the analysis, five of which were crossed over to the control arm after declining to receive HBOT. The other four required coronary artery bypass grafting or did not have a lesion suitable for stenting, while there were also four subjects excluded from the control group for the same reasons. [Dotsenko 2007](#) reported seven individuals in each group who withdrew early in the treatment phase and at final follow-up had lost a total of 14 participants (11%). Sensitivity analysis in this review has made best and worse case analyses (with the assumption of equal distribution of withdrawals) to examine potentially important effects on outcome where these studies contributed patients.

Intention-to-treat analysis

None of the included studies specifically indicated an intention to treat approach, and such an approach was not possible for [Sharifi 2004](#) as five subjects crossed from HBOT to control for analysis. [Swift 1992](#) reported full follow-up and did not report any protocol violation.

Effects of interventions

See: [Summary of findings for the main comparison hyperbaric oxygen therapy for acute coronary syndrome](#); [Summary of findings 2 hyperbaric oxygen therapy for acute coronary syndrome](#)

Primary outcomes


I. Death at any time after enrolment (Analysis I.1)

Figure 2

Five trials reported this outcome ([Dekleva 2004](#); [Dotsenko 2007](#); [Hot MI 1998](#); [Sharifi 2004](#); [Thurston 1973](#)), involving 614 subjects (92% of the total subjects in this review), with 287 (47%) allocated to standard treatment plus HBOT and 327 (53%) to standard therapy alone. [Thurston 1973](#) contributed 36% of the subjects to this analysis: 21 subjects (7.3%) died in the HBOT group versus 38 (11.6%) in the control group. There was a statistically significant reduction in the risk of death following HBOT (the RR of death with HBOT was 0.58, 95% CI 0.36 to 0.92, $P = 0.02$). This result, was however sensitive to the allocation of withdrawals (best case RR of death with HBOT is 0.37, 95% CI 0.23 to 0.58, $P < 0.001$, worst case RR 1.34, 95% CI 0.91 to 1.96, $P = 0.14$). Subgroup analysis suggested no significant difference between treatment groups for those in cardiogenic shock (RR with cardiogenic shock 0.61 95% CI 0.32 to 1.18, $P = 0.15$, but a benefit in those without cardiogenic shock (RR 0.57, 95% CI 0.33 to 0.98, $P = 0.004$). There was no indication of significant heterogeneity between trials ($I^2 = 0\%$). Despite the result

of the pooled analysis, the absolute risk difference (by simple χ^2 analysis) of 4.3% between control and HBOT is not statistically significant ($P = 0.07$), with an NNT to avoid one extra death of 24, (95% CI 259 in favour of control to 12 in favour of HBOT).

Figure 2. Forest plot of comparison: I Death, outcome: I.I Death at any time.



2. Major Adverse Coronary Events (MACE) (Analysis 2.1)

Only one trial reported this outcome (Sharifi 2004), involving 61 patients (9% of the total subjects in this review), with 24 (39%) analysed as receiving standard therapy with HBOT, and 37 (61%) standard therapy alone. One subject (4.2%) suffered with a MACE following HBOT versus 13 subjects (35.1%) in the control group. There was a statistically significant reduction in the risk of MACE following HBOT (RR 0.12, 95% CI 0.02 to 0.85, $P = 0.03$). This result was however, sensitive to the allocation of withdrawals (best case RR of death with HBOT is 0.09, 95% CI 0.01 to 0.61, $P = 0.01$, worst case RR 0.56, 95% CI 0.23 to 1.40, $P = 0.22$). The absolute risk difference of 30.9% between sham and HBOT is significant ($P = 0.005$), with an NNT to avoid one extra MACE of 4, (95% CI 3 to 10).

One trial reported the incidence of recurrent acute myocardial infarction (RAMI) at two years rather than MACE (Dotsenko 2007), and it was not clear if this included all those who died during the study period. We have therefore reported this outcome separately. This trial included 129 patients (19% of the total subjects in this

review), with 65 (50.4%) allocated to control and 64 (49.6%) to HBOT. Three subjects (4.7%) suffered with a RAMI following HBOT versus 11 subjects (16.9%) in the control group. There was a statistically significant reduction in the risk of RAMI following HBOT (RR 0.28, 95% CI 0.08 to 0.95, $P = 0.04$). This result was however, sensitive to the allocation of withdrawals (best case RR of death with HBOT is 0.17, 95% CI 0.05 to 0.55, $P = 0.0003$, worst case RR 0.92, 95% CI 0.42 to 2.02, $P = 0.84$). The absolute risk difference of 12.2% between sham and HBOT is significant ($P = 0.03$), with an NNT to avoid one extra MACE of 8, (95% CI 4 to 61).

Secondary outcomes

3. Significant cardiac events (only significant dysrhythmias were reported) (Analysis 3.1)

Only one trial reported this outcome (Thurston 1973) involving 208 patients (31% of the total subjects in this review), with 103

randomised to receive HBOT and 105 in the control arm. Of the events recorded, three dysrhythmias were accepted as 'significant cardiac events' - complete heart block, ventricular fibrillation and asystole. It is not clear if the numbers reported reflect individuals who suffered these events, or the number of events in total. Overall there were 25 such events reported in the patients receiving HBOT versus 43 such events in the control group, and patients receiving HBOT were significantly less likely to suffer one of these dysrhythmias (RR 0.59, 95% CI 0.39 to 0.89, $P = 0.01$). The absolute risk reduction of 17% corresponds to an NNT to avoid one event of 6, 95% CI 3 to 24. This result was however sensitive to the allocation of withdrawals, best case RR of significant event with HBOT is 0.51, 95%CI 0.34 to 0.77, $P = 0.001$, worst case RR 0.73, 95%CI 0.50 to 1.06, $P = 0.10$).

Separate analyses for each of the three dysrhythmias suggested HBOT patients were significantly less likely to suffer with complete heart block (RR 0.32, 95% CI 0.12 to 0.84, $P = 0.02$), but not ventricular fibrillation (RR 0.78, 95% CI 0.36 to 1.71, $P = 0.54$) or asystole (RR 0.73, 95% CI 0.34 to 1.56, $P = 0.42$).

4. Time to relief of cardiac pain (Analysis 4.1)

Only one trial contributed results to this outcome (Hot MI 1998) involving 81 subjects (12% of the total), 40 randomised to HBOT and 41 to control: 57 other subjects enrolled did not contribute data to this analysis, our best estimate is that these were 29 in the standard care plus HBOT group and 28 receiving standard care alone. The mean time to pain relief in the HBOT group was 261 minutes versus 614 minutes in the control group and this difference was statistically significant (WMD 353 minutes, 95% CI 219 to 488, $P < 0.0001$).

5. Size of infarct area

No trial reported any data on this outcome.

6. Magnitude of cardiac enzyme rise (Analysis 5.2, 5.2, 5.3)

Two trials contributed results to this outcome (Dekleva 2004 (as reported in Vlahovic 2004); Hot MI 1998) involving 184 subjects (28% of the total), 90 randomised to HBOT and 94 to control: 28 other subjects enrolled in Hot MI 1998 did not contribute data to this analysis, 16 in the standard care plus HBOT group and 12 receiving standard care alone. The Hot MI 1998 study reported serum creatine phosphokinase (CPK) levels at 12-hours post-treatment and 24-hours post-treatment, and both Hot MI 1998 and Dekleva 2004 reported the maximum level recorded. The levels at 12 and 24 hours were lower in the Hot MI 1998 patients receiving HBOT, but not statistically significantly so (12hrs- MD 138 international units [IU] lower with HBOT, 95% CI 843 lower to 568 higher, $P = 0.70$; 24 hrs MD 65 IU lower with HBOT, 95% CI 531lower to 401higher, $P = 0.78$). The maximum level

recorded was significantly lower following HBOT (MD 493 IU lower, 95% CI 839 to 148, $P = 0.005$).

7. Left ventricular function (Analysis 6.1 and 6.2)

Three trials reported on improvements in LV function (Dekleva 2004; Hot MI 1998; Swift 1992). Swift 1992 reported the number of individuals in whom improved function could be demonstrated on echocardiography following HBOT, while the other two reported LV ejection fraction (LVEF) at discharge (Hot MI 1998), day two and three weeks (Dekleva 2004). Dekleva 2004 also reported several other measures of cardiac function. Swift 1992 involved 34 subjects (5% of the total), 24 randomised to HBOT and 10 to control. 12 subjects showed improved contraction in at least one segment in the HBOT group versus zero in the control group. This difference was not, however, statistically significant (RR of improvement without HBOT 0.09, 95% CI 0.01 to 1.4, $P = 0.09$). Hot MI 1998 and Dekleva 2004 together involved 190 subjects (29% of the total), 94 randomised to HBOT and 96 to control. LVEF was significantly improved in those patients who received HBOT (MD 5.5% better, 95% CI 2.2% to 8.8%, $P = 0.001$). One control patient in Dekleva 2004 did not contribute to the analysis, while 21 subjects enrolled in the Hot MI 1998 study did not contribute data either, 12 in the HBOT group and 9 in the control.

8. Length of stay (Analysis 7.1)

Participants who were given HBOT had a mean stay in hospital of 7.4 days versus 9.2 days for those receiving the control treatment. This difference was not statistically significant (WMD 1.8 days, 95% CI 3.7 to -0.1, $P = 0.06$). Data were from 64 participants in the pilot phase of the Hot MI 1998 study (31 randomised to HBOT and 33 to control) 18 other participants did not contribute data to this analysis (10 from the HBOT arm and 8 from the control arm).

9. Myocardial perfusion

No trials reported any data on this outcome.

10. Quality of life

No trials reported any data on this outcome.

11. Rate of readmission

No trials reported any data on this outcome.

12. Costs of treatment

No trials attempted to estimate the cost-effectiveness of therapy.

Adverse effects

13. Tympanic membrane rupture (TMR), neurological oxygen toxicity and claustrophobia (Analysis 8.1)

Two trials reported on the incidence of tympanic membrane rupture due to barotrauma ([Sharifi 2004](#); [Thurston 1973](#)) involving 269 subjects (41% of the total), 127 (47%) randomised to HBOT and 142 (53%) randomised to control. One subject suffered TMR in the HBOT group versus none of the controls. This difference was not statistically significant (RR of TMR with HBOT 4.56, 95% CI 0.19 to 107.54, $P = 0.35$).

Three trials ([Hot MI 1998](#) (pilot phase); [Sharifi 2004](#); [Thurston 1973](#)) involving 274 subjects reported a zero incidence of neurological oxygen toxicity in all arms. No trial reported on any adverse effects in relation to standard therapeutic measures.

One trial reported on claustrophobia ([Thurston 1973](#)) involving 208 subjects (31% of the total), 103 (50%) randomised to HBOT and 105 (50%) to control. There were 15 subjects (15%) with claustrophobia requiring cessation of therapy in the HBOT group versus none in the control group. This difference is statistically significant (RR of claustrophobia with HBOT 31.6, 95% CI 1.92 to 521, $P = 0.02$).

ADDITIONAL SUMMARY OF FINDINGS *[Explanation]*

hyperbaric oxygen therapy for acute coronary syndrome						
Patient or population: patients with acute coronary syndrome Settings: acute care hospital Intervention: hyperbaric oxygen therapy						
Outcomes	Illustrative comparative risks* (95% CI)		Relative effect (95% CI)	No of Participants (studies)	Quality of the evidence (GRADE)	Comments
	Assumed risk	Corresponding risk				
	Control	hyperbaric oxygen therapy				
12 hour Plasma Creatine Phosphokinase		The mean 12 hour Plasma Creatine Phosphokinase in the intervention groups was 138 lower (843.83 lower to 567.83 higher)		84 (1 study)	⊕⊕⊕○ moderate ¹	
24 hour Plasma Creatine Phosphate		The mean 24 hour Plasma Creatine Phosphate in the intervention groups was 65 lower (530.96 lower to 400.96 higher)		72 (1 study)	⊕⊕⊕○ moderate ¹	
Maximum Plasma Creatine Phosphate		The mean Maximum Plasma Creatine Phosphate in the intervention groups was 493.16 lower (838.74 to 147.58 lower)		184 (2 studies)	⊕⊕⊕⊕ high	

*The basis for the **assumed risk** (e.g. the median control group risk across studies) is provided in footnotes. The **corresponding risk** (and its 95% confidence interval) is based on the assumed risk in the comparison group and the **relative effect** of the intervention (and its 95% CI).

CI: Confidence interval;

GRADE Working Group grades of evidence

High quality: Further research is very unlikely to change our confidence in the estimate of effect.

Moderate quality: Further research is likely to have an important impact on our confidence in the estimate of effect and may change the estimate.

Low quality: Further research is very likely to have an important impact on our confidence in the estimate of effect and is likely to change the estimate.

Very low quality: We are very uncertain about the estimate.

¹ Single trial only

DISCUSSION

Summary of main results

This review has included data from six trials investigating the treatment of ACS with HBOT, and we believe these represent all randomised human trials in this area, both published and unpublished, at the time of searching. One trial included subjects with ACS, while five included only subjects with confirmed myocardial infarction.

Pooled data for clinical outcomes of interest were performed with respect to the risk of death, cardiac enzyme peak levels, left ventricular function and adverse effects. CPK rose significantly less following HBOT (MD 493 IU, $P = 0.005$), implying a smaller volume of infarction, and LVEF was better following HBOT (MD 5.5%, $P = 0.001$). The clinical and functional significance of these differences is not clear. The risk of dying was significantly better following HBOT (RR 0.58, $P = 0.02$) and the absolute risk difference of 4.3% suggested an NNT of around 24 patients in order to avoid one death by the addition of HBOT. Only one trial (Thurston 1973) reported the fate of those presenting in cardiogenic shock, and while there was no statistically significant difference between groups in this small sample, it is worth noting that all survivors were from the HBOT group (three from seven subjects versus none from five). The one small study that reported MACE rather than death alone (Sharifi 2004) also suggested better outcome with the use of HBOT (RR 0.12, $P = 0.03$) with a risk difference of 31% and an NNT of 4. Similarly, Dotsenko 2007 reported a significant reduction in the chance of re-infarction with the administration of HBOT (RR 0.28, $P = 0.04$) with a risk difference of 12.2% and an NNT of 8. These possible treatment effects would be of great clinical importance and deserve further investigation.

Overall completeness and applicability of evidence

Only six trials with 665 participants were available for evaluation using our planned comparisons, and meta-analysis was not appropriate or possible for a number of these. The evidence is therefore very incomplete and cannot be applied directly to routine clinical care for patients with acute coronary syndrome.

These trials were published over a 24-year period up to 2007, and from a wide geographical area. We had planned to perform subgroup analyses with respect to inclusion or otherwise of thrombolysis, the nature of comparator treatments, dose of oxygen received (pressure, time and length of treatment course), the presence or absence of cardiac failure, the site of infarct, and to compare those with established versus pre-infarct subjects. However, the paucity of eligible trials and poor reporting suggested these analyses would not be informative. Patient inclusion criteria were not standard, and poorly reported in some trials. Only Hot MI

1998, Swift 1992 and Dotsenko 2007 clearly indicated the time at which the inclusion criteria were applied. There was significant variation both in oxygen dose during an individual treatment session, and in the number of sessions administered to each patient. While all trials used some form of 'standard' cardiac therapy in a dedicated unit designed to maximise outcome, these comparator therapies were generally poorly described and could not form the basis for a meaningful subgroup analysis.

Quality of the evidence

As well as carrying little statistical power, other problems for this review were the variable methodological quality of many of these trials, differences in entry criteria and the nature and timing of outcomes, and poor reporting of both outcomes and methodology. In particular, there is a possibility of bias due to different anatomical locations and extent of myocardial damage on entry to these small trials, as well as from non-blinded management decisions in all except Swift 1992. Further, we could only pool the data for two of our outcomes of interest given little commonality between the outcomes reported in these trials.

For the primary outcomes, only a single trial clearly recorded all major adverse copronary events (MACE) and we were unable to calculate a pooled estimate of the effect of allocation to receive HBOT. For the risk of death following acute coronary syndrome, our estimate is that patients allocated to HBOT were associated with a reduction in the risk of death by around 42% (RR: 0.58, 95% CI 0.36 to 0.92). This estimate was derived from only five trials including 614 subjects and the GRADE approach rates this evidence as of low quality following downgrading because this critical outcome had a low incidence in this small group of patients. For our secondary outcomes, we could pool only the maximum plasma creatine phosphate (CPK) for two trials. Using the GRADE approach, the evidence for lower CPK associated with patients who received HBOT was rated as high.

As is common with small trials, the incidence of adverse effects was poorly assessed by the studies included in this review. No trial reported any neurological or pulmonary oxygen toxicity in any group, while there was only one reported case of severe ear barotrauma as a consequence of compression. Thurston 1973 reported 15 individuals who needed to be removed from a single occupancy hyperbaric chamber because of claustrophobia, a rate of 15%. While this is a clinically significant problem in that trial, it is unlikely this rate would be sustained when using larger compression chambers designed for multiple occupancy. There are a number of more minor complications that may occur commonly. Visual disturbance, usually reduction in visual acuity secondary to conformational changes in the lens, is very commonly reported - perhaps as many as 50% of those having a course of 30 treatments (Khan 2003). While the great majority of patients recover spontaneously over a period of days to weeks, a small proportion of patients continue to require correction to restore sight to pre-

treatment levels. None of the trials included in this review reported visual changes. The second most common adverse effect associated with HBOT is middle-ear barotrauma. Barotrauma can affect any air-filled cavity in the body (including the middle ear, lungs and respiratory sinuses) and occurs as a direct result of compression. Ear barotrauma is by far the most common as the middle ear air space is small, largely surrounded by bone and the sensitive tympanic membrane, and usually requires active effort by the patient in order to inflate the middle ear through the eustachian tube on each side. Barotrauma is thus not a consequence of HBOT directly, but rather of the physical conditions required to administer it. Most episodes of barotrauma are mild, easily treated or recover spontaneously and do not require the therapy to be abandoned.

Potential biases in the review process

The authors of this review have no conflict of interest to declare and none are authors of any literature contributing to this review. We believe we have conducted this review without bias. All of these findings are, however, subject to a potential publication bias. While we have made every effort to locate further unpublished data, it remains possible that this review is subject to a positive publication bias, with generally favourable trials more likely to achieve reporting. With regard to any effect on the quality of life for these patients, we have located no relevant data.

Agreements and disagreements with other studies or reviews

Although there has been relatively little clinical literature on this subject, these encouraging findings are in general agreement with opinion in the literature [Ellestad 2009](#). We are not aware of any other formal systematic reviews of the literature.

AUTHORS' CONCLUSIONS

Implications for practice

There is some evidence from this review that HBOT following an episode of ACS reduces the risk of death, re-infarction, significant dysrhythmias and MACE, as well as limiting the peak levels of CPK and improving left ventricular ejection fraction. HBOT may also reduce the time required to achieve relief from cardiac ischaemic pain in these patients. The small number of studies, the

modest numbers of patients, and the methodological and reporting inadequacies of the primary studies included in this review demand a cautious interpretation. Thus, the routine adjunctive use of HBOT in these patients cannot be justified by this review.

Implications for research

Given the indicative findings of improved outcomes with the use of HBOT in these patients, there is a case for large randomised trials of high methodological rigour in order to define the true extent of benefit (if any) from the administration of HBOT. Specifically, more information is required on the subset of disease severity and timing of therapy most likely to result in benefit from this therapy. Given the activity of HBOT in modifying ischaemia-reperfusion injury, attention should be given to combinations of HBOT and thrombolysis in the early treatment of acute coronary events and the prevention of re-stenosis after stent placement. Any future trials would need to consider in particular:

- appropriate sample sizes with power to detect the expected differences suggested by this review;
- careful definition and selection of target patients;
- acute versus sub-acute administration of HBOT;
- appropriate range of oxygen doses per treatment session (pressure and time);
- appropriate and carefully defined comparator therapy;
- use of an effective sham therapy;
- effective and explicit blinding of outcome assessors;
- appropriate outcome measures including all those listed in this review;
- careful elucidation of any adverse effects;
- the cost-utility of the therapy;
- patient quality of life.

ACKNOWLEDGEMENTS

The authors wish to acknowledge the help of the Cochrane Heart Group, and in particular Margaret Burke for her assistance with the development of the search strategy, and both Theresa Moore and Katherine Wornell for their patient comments and suggestions.

REFERENCES

References to studies included in this review

Dekleva 2004 *{published data only}*

* Dekleva M, Neskovic A, Vlahovic A, Putnikovic B, Beleslin B, Ostojic M. Adjunctive effect of hyperbaric oxygen treatment after thrombolysis on left ventricular function in patients with acute myocardial infarction. *American Heart Journal* 2004;**148**:e14.
Vlahovic A, Neskovic AN, Dekleva M, Putnikovic B, Popovic ZB, Otasevic P, et al. Hyperbaric oxygen treatment does not affect left ventricular chamber stiffness after myocardial infarction treated with thrombolysis. *American Heart Journal* 2004;**148**:e1.

Dotsenko 2007 *{published data only}*

Aliprandi-Costa B, Ranasinghe I, Chow V, Kapila S, Juergens C, Devlin G, et al. Management and outcomes of patients with acute coronary syndromes in Australia and New Zealand, 2000-2007. *MJA* 2011;**195**(3):116-21.
* Dotsenko EA, Salivonchik OP, Kozyro VI. Long term results of the use of hyperbaric oxygenation in patients with acute myocardial infarction. *Kardiologija* 2007;**12**:53-6.
Dotsenko EA, Salivonchik OP, Nikulina N, Welcome MO. The influence of hyperbaric oxygenation therapy on recurrent myocardial infarction and two-year survival rate in acute myocardial infarction patients. *Port Harcourt Medical Journal* 2009;**3**(3):256-63.

Hot MI 1998 *{published data only}*

Dekleva MN, Ostojic M, Vujnovic D. Hyperbaric oxygen and thrombolysis in acute myocardial infarction: a preliminary report. In: Sitinen SA, Leinio M editor(s). *Proceedings of the Twenty-first Annual Meeting of the European Underwater and Baromedical Society (EUBS)*. Helsinki: EUBS, 1995:9-13.
Shandling AH, Ellestad MH, Hart GB, Crump R, Marlow D, Van Natta B, et al. Hyperbaric oxygen and thrombolysis in myocardial infarction: The "HOT MI" pilot study. *American Heart Journal* 1997;**134**:544-50.
* Stavitsky Y, Shandling AH, Ellestad MH, Hart GB, Van Natta B, Messenger JC, et al. Hyperbaric oxygen and thrombolysis in myocardial infarction: the 'HOT MI' randomised multicenter study. *Cardiology* 1998;**90**:131-6.

Sharifi 2004 *{published data only}*

* Sharifi M, Fares W, Abdel-Karim I, Koch JM, Sopko J, Adler D. Usefulness of hyperbaric oxygen therapy to inhibit restenosis after percutaneous coronary intervention for acute myocardial infarction or unstable angina pectoris. *American Journal of Cardiology* 2004;**93**(June 15):1533-5.
Sharifi M, Fares W, Abdel-Karim I, Petrea D, Koch JM, Adler D, et al. Inhibition of restenosis by hyperbaric oxygen: a novel indication for an old modality. *Cardiovascular Radiation Medicine* 2002;**3**:124-6.

Swift 1992 *{published data only}*

Swift PC, Turner JH, Oxer HF, O'Sea JP, Lane GK, Woollard KV. Myocardial hibernation identified by hyperbaric oxygen treatment and echocardiography in

postinfarction patients: comparison with exercise thallium scintigraphy. *American Heart Journal* 1992;**124**:1151-8.

Thurston 1973 *{published data only}*

Thurston GJ, Greenwood TW, Bending MR, Connor H, Curwen MP. A controlled investigation into the effects of hyperbaric oxygen on mortality following acute myocardial infarction. *Quarterly Journal of Medicine* 1973;**XLII**:751-70.

References to studies excluded from this review

Cameron 1965 *{published data only}*

Cameron AJ, Gibb BH, Ledingham I. A controlled clinical trial of hyperbaric oxygen in the treatment of acute myocardial infarction. Hyperbaric Oxygenation: Proceedings of the Second International Congress. London: ES Livingstone, 1965:277.

Ciocatto 1965 *{published data only}*

Ciocatto E, Moricca G, Querci M, Cabrai M. Experimental studies on hyperbaric oxygenation. *Panminerva Medica* 1965;**7**(11):419-24.

Dai 1995 *{published data only}*

Dai R, Wu JT, Liu LP, Wang YQ. Hyperbaric oxygen for coronary heart disease. *Chinese Journal of Physical Therapy* 1995;**18**:236-7.

Markarian 1991 *{published data only}*

Markarian SS, Shirinskaia GI, Starostin SG, Zagvozkin VN. Comparative study of the effect of drug therapy and its combination with hyperbaric oxygenation and hemosorption on stable angina pectoris [Sravnitel'noe izuchenie vlianiia medikamentoznoi terapii i ee sochetanii s giperbaricheskoi oksigenatsiei i gemocorbtsei na stabil'nuu stenokardiiu]. *Kardiologija* 1991;**31**(9):40-2.

Thomas 1990 *{published data only}*

Thomas MP, Brown LA, Sponseller DR, Williamson SE, Diaz JA, Guyton DP. Myocardial infarct size reduction by the synergistic effect of hyperbaric oxygen and recombinant tissue plasminogen activator. *American Heart Journal* 1990;**120**(4):791-800.

Additional references

Ashfield 1969

Ashfield R, Drew CE, Gavey CJ. Severe acute myocardial infarction treated with hyperbaric oxygen. *Postgraduate Medical Journal* 1969;**45**:648-53.

Bennett 2002

Bennett. The database of randomised controlled trials in hyperbaric medicine. www.hboevidence.com (accessed on 21 February 2005).

BHF 2012

British Heart Foundation. Coronary heart disease statistics. <https://www.bhf.org.uk/publications/statistics/coronary-heart-disease-statistics-2012>. London: British Heart Foundation, 3/12/2012.

Boerema 1960

Boerema I, Meyne NG, Brummelkamp WK, Bouma S, Mensch MH, Kamermans F, et al. Life without blood: a study of the influence of high atmospheric pressure and hypothermia on dilution of blood. *Journal of Cardiovascular Surgery* 1960;1(1):133–46.

Butler 2008

Butler FK, Hagan C. Ocular complications in hyperbaric oxygen therapy. In: Neuman T, Thom S editor(s). *Physiology and Medicine of Hyperbaric Oxygen Therapy*. 1st Edition. Philadelphia: Saunders Elsevier, 2008:265–72.

Clark 2008

Clarke JM. Oxygen toxicity. In: Neuman T, Thom S editor(s). *Physiology and Medicine of Hyperbaric Oxygen Therapy*. 1st Edition. Philadelphia: Saunders Elsevier, 2008:527–564.

Ellestad 2009

Ellestad MH. Hyperbaric oxygen: its application in cardiology: a historical perspective and personal journey. *Cardiology Review* 2009;17(6):280–2.

Hammarlund 1999

Hammarlund C. The physiologic effects of hyperbaric oxygenation. In: Kindwall EP, Whelan HT editor(s). *Hyperbaric Medicine Practice*. 2nd Edition. Flagstaff Az: Best Publishing Company, 1999:37–68.

Harabin 1990

Harabin AL, Braisted JC, Flynn ET. Response of antioxidant enzymes to intermittent and continuous hyperbaric oxygen. *Journal of Applied Physiology* 1990;69(1):328–35.

Heistad 2003

Heistad D. Unstable coronary-artery plaques. *New England Journal of Medicine* 2003;349(24):2285–7.

Heyboer 2014

Heyboer M, Milovanova TN, Wojcik S, Grant W, Chin M, Hardy KR, et al. CD34+/CD45-dim stem cell mobilization by hyperbaric oxygen - changes with oxygen dosing. *Stem Cell Research* 2014;12(3):638–45.

Higgins 2008

Higgins JPT, Green S, editors. *Cochrane Handbook for Systematic Reviews of Interventions*. *Cochrane Book Series*. Chichester: Wiley-Blackwell, 2008.

Jain 2009

Jain KK. *Textbook of Hyperbaric Medicine*. 5th Edition. Seattle: Hogrefe and Huber, 2009.

Jones 2010

Jones SR, Carpin KM, Woodward SM, Khiabani KT, Stephenson LL, Wang WZ, et al. Hyperbaric oxygen Inhibits ischemia-reperfusion-induced neutrophil CD18Polarization by a nitric oxide mechanism. *Plastic and Reconstructive Surgery* 2010;126(2):403–11.

Kalla 2006

Kalla K, Christ G, Karnik R, Malzer R, Norman G, Prachar H, et al. Implementation of guidelines improves the standard of care the Viennese Registry on Reperfusion

Strategies in ST-Elevation Myocardial Infarction (Vienna STEMI Registry). *Circulation* 2006;113:2398–405.

Khan 2003

Khan B, Evans AW, Easterbrook M. Refractive changes in patients undergoing hyperbaric oxygen therapy: a prospective study. *Undersea and Hyperbaric Medicine* 2003;24:9.

Kindwall 2008

Kindwall EP, Whelan HT. *Hyperbaric Medicine Practice*. 3rd Edition. Flagstaff: Best Publishing Company, 2008.

Kline 1970

Kline HJ, Marano AJ, Johnson CD, Goodman P, Jacobson JH, Kuhn LA. Hemodynamic and metabolic effects of hyperbaric oxygenation in myocardial infarction. *Journal of Applied Pathology* 1970;28:256–63.

Leach 1998

Leach RM, Rees PJ, Wilmshurst P. ABC of oxygen. Hyperbaric oxygen therapy. *BMJ* 1998;317:1140–3.

Mathieu 2006

Mathieu D. *Handbook on Hyperbaric Medicine*. 2nd Edition. Milan: Springer, 2006.

Moon 1964

Moon AJ, Williams KG, Hopkinson WI. A patient with coronary thrombosis treated with hyperbaric oxygen. *Lancet* 1964;1:18–20.

Naghavi 2003

Naghavi M, Libby P, Falk E, Casscells SW, Litovsky S, et al. From vulnerable plaque to vulnerable patient: a call for new definitions and risk assessment strategies: part I. *Circulation* 2003;108(14):1664–72.

Nichols 2012

Nichols M, Townsend N, Luengo-Fernandez R, Leal J, Grey A, Scarborough P, Rayner M. European Cardiovascular Disease Statistics 2012. *European Heart Network, Brussels, European Society of Cardiology, Sophia Antipolis*. Brussels: European Society of Cardiology, Sophia Antipolis, 2012.

Sheffield 1988

Sheffield P. Tissue oxygen measurements. *Problem Wounds. The Role of Oxygen*. 1st Edition. New York: Elsevier Science Publishing, 1988:17–51.

Shupak 2008

Shupak A, Gilbey P. Effects of pressure. In: Neuman T, Thom S editor(s). *Physiology and Medicine of Hyperbaric Oxygen Therapy*. 1st Edition. Philadelphia: Saunders Elsevier, 2008:513–26.

Smith 1958

Smith G, Lawson DA. Experimental coronary arterial occlusion: effects of the administration of oxygen under pressure. *Scottish Medical Journal* 1958;3:346–50.

Speit 2000

Speit G, Dennog C, Eichorn U, Rothfuss A, Kaina B. Induction of heme oxygenase-1 and adaptive protection against the induction of DNA damage after hyperbaric oxygen treatment. *Cardiogenesis* 2000;21(10):1795–9.

Thom 1991

Thom SR, Elbuku M. Oxygen-dependent antagonism of lipid peroxidation in the rat. *Free Radical Biology and Medicine* 1991;**10**(6):413–26.

Thom 2009

Thom S. Oxidative stress is fundamental to hyperbaric oxygen therapy. *Journal of Applied Physiology* 2009;**106**: 988–95.

Tjarnstrom 1999

Tjarnstrom J, Wikstrom T, Bagge U. Effects of hyperbaric oxygen treatment on neutrophil activation and pulmonary sequestration in intestinal ischemia-reperfusion in rats. *European Surgical Research* 1999;**31**(2):147–54.

Wells 1977

Wells CH, Goodpasture JE, Horrigan DJ. Tissue gas measurements during hyperbaric oxygen exposure. Proceedings of the Sixth International Congress on Hyperbaric Medicine. Aberdeen: Aberdeen University Press, 1977:118–24.

WHO 2013

World Health Organisation. Cardiovascular diseases (CVDs). Fact sheet N°317 Updated March 2013. <http://www.who.int/mediacentre/factsheets/fs317/en/> (accessed on 19 December 2014).

* Indicates the major publication for the study

CHARACTERISTICS OF STUDIES

Characteristics of included studies [ordered by study ID]

Dekleva 2004

Methods	Unblinded randomised controlled trial, method of allocation and randomisation not described
Participants	74 subjects enrolled with firm diagnosis of AMI. Excluded those with heart failure, severe arrhythmias and over 70 years. The mean age of subjects in the HBOT group was 55 yrs (sd 7) and in the control group 54 yrs (sd 8). 22% of the HBOT group were female, as were 8% of the control group and 22% were diabetic compared to 5% of the controls
Interventions	Control had thrombolysis with 1.5m IU streptokinase over 30 minutes. Experimental group had the same, plus a single session of HBOT at 2.0 ATA for 60 minutes (mean time to treatment 13hrs)
Outcomes	Peak creatine kinase, LV function, death
Notes	Echocardiographer was blinded to therapy.

Risk of bias

Bias	Authors' judgement	Support for judgement
Random sequence generation (selection bias)	Low risk	"With a random number table, patients were randomly assigned to.."
Allocation concealment (selection bias)	Unclear risk	No description of attempts at concealment.
Blinding (performance bias and detection bias) All outcomes	High risk	There was no sham therapy and the patient and investigators were all aware of group allocation. "The patients randomly assigned to streptokinase plus HBO were transferred to the hyperbaric unit in the first 24 hours from the onset of symptoms and after thrombolytic therapy." Echocardiographer was blinded to therapy
Incomplete outcome data (attrition bias) All outcomes	Low risk	No withdrawals or loss to follow-up.
Selective reporting (reporting bias)	Low risk	The published report likely contains all relevant outcomes intended
Other bias	Low risk	No clear source suggesting other biases.

Dotsenko 2007

Methods	Unblinded randomised controlled trial. Randomisation by computer generated sequence, but allocation method unclear
Participants	129 subjects enrolled with ECG or biochemical evidence of AMI between 3 and 10 days prior to enrolment and aged 30 to 75 years. The mean age of subjects in both the HBOT and control groups was 55 yrs (sd 1). The authors stated that the sex ratio was the same for both groups but did not give the exact figures
Interventions	Control group had usual therapy including thrombolysis when indicated. Experimental group the same with the addition of HBOT at 1.3ATA for 40 minutes daily for six days
Outcomes	Mortality, reinfarction
Notes	Not possible to tell if the two outcomes are mutually exclusive or not

Risk of bias

Bias	Authors' judgement	Support for judgement
Random sequence generation (selection bias)	Low risk	Computer generated random number table ("randomly divided into two groups with computer technology")
Allocation concealment (selection bias)	Unclear risk	No statement in report.
Blinding (performance bias and detection bias) All outcomes	High risk	No sham therapy administered.
Incomplete outcome data (attrition bias) All outcomes	Low risk	Seven participants in each arm refused to continue in the study shortly after enrolment
Selective reporting (reporting bias)	Low risk	No evidence of outcomes not reported
Other bias	Unclear risk	Little information upon which to make this judgement.

Hot MI 1998

Methods	Multicentred, randomised trial. Allocation method not described. No blinding. 16 subjects excluded after randomisation
Participants	138 subjects enrolled in emergency room the numbers randomised to each arm were not reported. Patients in emergency room with AMI diagnosed by clinical features and ECG changes, and who were eligible for thrombolysis. Age 18 to 80 years. 16 excluded due to haemodynamic instability, no proven AMI, exceeded time limit for thrombolysis, incorrect protocol, incomplete data or refusal of HBOT. The mean age of subjects in

Hot MI 1998 (Continued)

	both the HBOT and control groups was 59 yrs (sd 12) . 19% of the HBOT group were female, as were 26% of the control group
Interventions	Controls received thrombolysis, aspirin, heparin and intravenous nitroglycerine. HBO group received the same plus 1 treatment of 2ATA 100% oxygen for 2 hours
Outcomes	Death, time to pain relief, magnitude of enzyme change, left ventricular ejection fraction. Length of stay
Notes	

Risk of bias

Bias	Authors' judgement	Support for judgement
Random sequence generation (selection bias)	Low risk	"With a random number table, patients were randomised to ..."
Allocation concealment (selection bias)	Unclear risk	No description given of possible allocation concealment.
Blinding (performance bias and detection bias) All outcomes	High risk	There was no sham therapy. "The patients randomised to HBO were then immediately transferred to the hyperbaric unit..". LVEF measures were observer blinded
Incomplete outcome data (attrition bias) All outcomes	Low risk	Missing data balanced between groups and unlikely to have affected result. Exclusions described
Selective reporting (reporting bias)	Low risk	All expected outcomes seem to be included.
Other bias	High risk	No indication of other important sources of bias.

Sharifi 2004

Methods	Randomised controlled trial without blinding or allocation concealment. Patients refusing HBOT crossed over to control (5 subjects). Analysis by intention to treat is therefore not possible
Participants	69 subjects enrolled (33 HBOT, 36 control) with clinical diagnosis of acute AMI or unstable angina, but were excluded if pain was ongoing, or S-T segments unresolved after 30 minutes of medical therapy. The mean age of subjects in the HBOT group was 63 yrs (sd 12) and in the control group 65 yrs (sd 13). 42% of the HBOT group were female, as were 43% of the control group 5 subjects crossed from HBOT to control after refusal or early termination of HBOT, while a further 4 subjects from each group did not require PCI. Therefore final analysis of 24 HBOT and 37 control subjects

Interventions	Controls underwent stenting and received aspirin, heparin and clopidogrel. Experimental subjects received HBOT at 2ATA for 90 minutes 1 hour prior to or immediately following stent, and a second treatment within 18 hours. Medical therapy was the same for both groups	
Outcomes	MACE, adverse events	
Notes		
<i>Risk of bias</i>		
Bias	Authors' judgement	Support for judgement
Random sequence generation (selection bias)	Unclear risk	No method of randomisation was described. "33 were randomised to the HOT arm and 36 to the control group..."
Allocation concealment (selection bias)	Unclear risk	No description of allocation concealment.
Blinding (performance bias and detection bias) All outcomes	High risk	There was no sham therapy. "All patients in the HOT arm underwent two hyperbaric dives..."
Incomplete outcome data (attrition bias) All outcomes	High risk	Five patients crossed from HBOT to control arm. ITT analysis not possible
Selective reporting (reporting bias)	Low risk	"The prespecified primary endpoints consisted of the composite endpoints of death, MI, emergent coronary bypass surgery and target lesion revascularization...". All were reported
Other bias	Low risk	No indication of other significant bias.

Swift 1992

Methods	Randomised controlled trial with concealed allocation. Schedule called for 2 active for each control subject. No loss to follow-up and subjects were blinded with sham therapy
Participants	34 subjects (24 HBOT, 10 control) enrolled with firm clinical diagnosis of AMI within the past week, plus abnormal wall motion on transoesophageal echo. Uncontrolled heart failure excluded. Most had received thrombolysis. The mean age of subjects in the study was 58 yrs with a range from 27 to 70 yrs - the figures for each group were not given. 10% of the subjects were female

Swift 1992 (Continued)

Interventions	Control group had echocardiography, exposure to 2ATA breathing air for 30 minutes and repeat echo. HBOT group had same schedule but breathed 100% oxygen at 2ATA
Outcomes	Improved LV function on echocardiography. No follow-up past the immediate post-HBOT phase. Outcome assessors were blinded and shown results in random sequence
Notes	Perhaps not designed as a therapeutic trial, but does satisfy entry criteria and measured a short-term outcome

Risk of bias

Bias	Authors' judgement	Support for judgement
Random sequence generation (selection bias)	Unclear risk	No method of sequence allocation described. "Patients... were randomly allocated to received either room air or 100% oxygen."
Allocation concealment (selection bias)	Unclear risk	No method described.
Blinding (performance bias and detection bias) All outcomes	Low risk	Sham therapy such that patient and investigators were blind. "Patients were pressurised to 2ATA for 30 minutes and were randomly allocated...."
Incomplete outcome data (attrition bias) All outcomes	Unclear risk	Some results are composite outcomes of individual segments of heart, so hard to tell if there is missing data. All individuals seem to be represented
Selective reporting (reporting bias)	Low risk	All outcomes of interest to investigators seem to be reported
Other bias	Low risk	No obvious source of bias.

Thurston 1973

Methods	Sealed envelope randomisation, no blinding after allocation to group. 13 subjects withdrawn due to misdiagnosis or age recorded wrongly
Participants	221 subjects (110 HBOT, 111 control) with strong clinical probability of myocardial infarction at admission, aged <70 years. 13 later excluded because of misdiagnosis or exceeded age limit. The mean age of subjects in the two groups was not give, but the age distribution was similar in the two groups, with the majority of subjects aged between 45 and 64 years. 15% of the HBOT group were female, as were 17% of the control group

Thurston 1973 (Continued)

Interventions	Control: “full orthodox coronary care including oxygen at 6 lpm by mask.” HBOT: As above, minus mask oxygen and plus HBOT at 2ATA for 2 hours, followed by 1 hour on air at 1ATA, repeating for 48 hours	
Outcomes	Death at 3 weeks, rate of significant dysrhythmias, adverse effects. MACE not given as death and significant dysrhythmia may have been reported in the same individual	
Notes	Some indication that HBOT subjects may have been more severely ill than control. Quality assessment: Randomisation: not described, Allocation: B, Performance Bias: unblinded, Detection bias: not described	
<i>Risk of bias</i>		
Bias	Authors’ judgement	Support for judgement
Random sequence generation (selection bias)	Unclear risk	Method unclear. “...sealed envelopes giving the random allocation..”
Allocation concealment (selection bias)	Low risk	Used sealed envelopes to reveal allocation. “Peel Index (was done)...as soon as possible after entry into the trial and before opening the sealed envelopes giving the random allocation into treatment and control groups.”
Blinding (performance bias and detection bias) All outcomes	High risk	No sham therapy.
Incomplete outcome data (attrition bias) All outcomes	Low risk	Primary outcome accounted for all patients after exclusions.
Selective reporting (reporting bias)	Low risk	All outcomes of interest apparently addressed.
Other bias	Unclear risk	Despite randomisation, the HBOT group was in general a little more unwell

AMI - acute myocardial infarction
ATA - atmospheres absolute
HBOT - Hyperbaric Oxygen therapy
lpm - litres per minute

Characteristics of excluded studies *[ordered by study ID]*

Study	Reason for exclusion
Cameron 1965	Case series, no comparator group
Ciocatto 1965	This is an animal experiment. "The experiments were conducted on unselected rabbits"
Dai 1995	Not an RCT. "Based on financial situations, the participants were separated into 2 groups"
Markarian 1991	RCT enrolling patients with angina, including unstable angina, but cannot obtain results broken down by functional class
Thomas 1990	Animal study

DATA AND ANALYSES

Comparison 1. Death

Outcome or subgroup title	No. of studies	No. of participants	Statistical method	Effect size
1 Death at any time	5	614	Risk Ratio (M-H, Fixed, 95% CI)	0.58 [0.36, 0.92]
1.1 Subjects presenting in cardiogenic shock	1	12	Risk Ratio (M-H, Fixed, 95% CI)	0.61 [0.32, 1.18]
1.2 Subjects presenting without cardiogenic shock	5	602	Risk Ratio (M-H, Fixed, 95% CI)	0.57 [0.33, 0.98]
2 Death - best case scenario	5	617	Risk Ratio (M-H, Fixed, 95% CI)	0.37 [0.23, 0.58]
3 Death - worst case scenario	5	617	Risk Ratio (M-H, Fixed, 95% CI)	1.34 [0.91, 1.96]

Comparison 2. Major Adverse Cardiac Events

Outcome or subgroup title	No. of studies	No. of participants	Statistical method	Effect size
1 Major Adverse Cardiac Events	1	61	Risk Ratio (M-H, Fixed, 95% CI)	0.12 [0.02, 0.85]
2 MACE - Best case scenario	1	69	Risk Ratio (M-H, Fixed, 95% CI)	0.09 [0.01, 0.61]
3 MACE - worst case scenario	1	69	Risk Ratio (M-H, Fixed, 95% CI)	0.56 [0.23, 1.40]
4 Recurrent acute myocardial infarction (AMI)	1	129	Risk Ratio (M-H, Fixed, 95% CI)	0.28 [0.08, 0.95]
5 Recurrent AMI - best case scenario	1	129	Risk Ratio (M-H, Fixed, 95% CI)	0.17 [0.05, 0.55]
6 Recurrent AMI - Worst case scenario	1	129	Risk Ratio (M-H, Fixed, 95% CI)	0.92 [0.42, 2.02]

Comparison 3. Significant dysrhythmias (complete heart block, ventricular fibrillation, asystole)

Outcome or subgroup title	No. of studies	No. of participants	Statistical method	Effect size
1 Overall (CHB, VF and asystole combined)	1	208	Risk Ratio (M-H, Fixed, 95% CI)	0.59 [0.39, 0.89]
2 Significant dysrhythmias (complete heart block, ventricular fibrillation or asystole)	1		Risk Ratio (M-H, Fixed, 95% CI)	Subtotals only
2.1 Complete heart block	1	208	Risk Ratio (M-H, Fixed, 95% CI)	0.32 [0.12, 0.84]
2.2 Ventricular fibrillation	1	208	Risk Ratio (M-H, Fixed, 95% CI)	0.78 [0.36, 1.71]
2.3 Asystole	1	208	Risk Ratio (M-H, Fixed, 95% CI)	0.73 [0.34, 1.56]
3 Overall best case	1	221	Risk Ratio (M-H, Fixed, 95% CI)	0.51 [0.34, 0.77]

4 Overall worst case	1	221	Risk Ratio (M-H, Fixed, 95% CI)	0.73 [0.50, 1.06]
----------------------	---	-----	---------------------------------	-------------------

Comparison 4. Time to pain relief

Outcome or subgroup title	No. of studies	No. of participants	Statistical method	Effect size
1 Time to relief of pain	1	81	Mean Difference (IV, Fixed, 95% CI)	-353.00 [-487.55, -218.45]

Comparison 5. Magnitude of cardiac enzyme changes

Outcome or subgroup title	No. of studies	No. of participants	Statistical method	Effect size
1 12 hour Plasma Creatine Phosphokinase	1	84	Mean Difference (IV, Fixed, 95% CI)	-138.0 [-843.83, 567.83]
2 24 hour Plasma Creatine Phosphate	1	72	Mean Difference (IV, Fixed, 95% CI)	-65.0 [-530.96, 400.96]
3 Maximum Plasma Creatine Phosphate	2	184	Mean Difference (IV, Fixed, 95% CI)	-493.16 [-838.74, -147.58]

Comparison 6. Improvement in left ventricular function

Outcome or subgroup title	No. of studies	No. of participants	Statistical method	Effect size
1 Improved contraction in at least one segment (post-HBOT echo)	1	34	Risk Ratio (M-H, Fixed, 95% CI)	0.09 [0.01, 1.40]
2 Left Ventricular Ejection Fraction- % (final estimate)	2	190	Mean Difference (IV, Random, 95% CI)	5.47 [2.19, 8.75]

Comparison 7. Length of Stay

Outcome or subgroup title	No. of studies	No. of participants	Statistical method	Effect size
1 Overall length of stay (days)	1	64	Mean Difference (IV, Fixed, 95% CI)	-1.80 [-3.70, 0.10]

Comparison 8. Adverse events of therapy


Outcome or subgroup title	No. of studies	No. of participants	Statistical method	Effect size
1 Total adverse events	3		Risk Ratio (M-H, Fixed, 95% CI)	Subtotals only
1.1 Tympanic membrane rupture	2	269	Risk Ratio (M-H, Fixed, 95% CI)	4.56 [0.19, 107.54]
1.2 Acute neurological oxygen toxicity	2	274	Risk Ratio (M-H, Fixed, 95% CI)	0.0 [0.0, 0.0]
1.3 Claustrophobia	1	208	Risk Ratio (M-H, Fixed, 95% CI)	31.60 [1.92, 521.22]

Analysis 1.1. Comparison 1 Death, Outcome 1 Death at any time.


Review: Hyperbaric oxygen therapy for acute coronary syndrome

Comparison: 1 Death

Outcome: 1 Death at any time



(Continued ...)




Analysis 1.2. Comparison 1 Death, Outcome 2 Death - best case scenario.

Review: Hyperbaric oxygen therapy for acute coronary syndrome

Comparison: 1 Death

Outcome: 2 Death - best case scenario




Analysis 1.3. Comparison 1 Death, Outcome 3 Death - worst case scenario.

Review: Hyperbaric oxygen therapy for acute coronary syndrome

Comparison: 1 Death

Outcome: 3 Death - worst case scenario




Analysis 2.1. Comparison 2 Major Adverse Cardiac Events, Outcome 1 Major Adverse Cardiac Events.

Review: Hyperbaric oxygen therapy for acute coronary syndrome

Comparison: 2 Major Adverse Cardiac Events

Outcome: 1 Major Adverse Cardiac Events




Analysis 2.2. Comparison 2 Major Adverse Cardiac Events, Outcome 2 MACE - Best case scenario.

Review: Hyperbaric oxygen therapy for acute coronary syndrome

Comparison: 2 Major Adverse Cardiac Events

Outcome: 2 MACE - Best case scenario




Analysis 2.3. Comparison 2 Major Adverse Cardiac Events, Outcome 3 MACE - worst case scenario.

Review: Hyperbaric oxygen therapy for acute coronary syndrome

Comparison: 2 Major Adverse Cardiac Events

Outcome: 3 MACE - worst case scenario




Analysis 2.4. Comparison 2 Major Adverse Cardiac Events, Outcome 4 Recurrent acute myocardial infarction (AMI).

Review: Hyperbaric oxygen therapy for acute coronary syndrome

Comparison: 2 Major Adverse Cardiac Events

Outcome: 4 Recurrent acute myocardial infarction (AMI)




Analysis 2.5. Comparison 2 Major Adverse Cardiac Events, Outcome 5 Recurrent AMI - best case scenario.

Review: Hyperbaric oxygen therapy for acute coronary syndrome

Comparison: 2 Major Adverse Cardiac Events

Outcome: 5 Recurrent AMI - best case scenario




Analysis 2.6. Comparison 2 Major Adverse Cardiac Events, Outcome 6 Recurrent AMI - Worst case scenario.

Review: Hyperbaric oxygen therapy for acute coronary syndrome

Comparison: 2 Major Adverse Cardiac Events

Outcome: 6 Recurrent AMI - Worst case scenario




Analysis 3.1. Comparison 3 Significant dysrhythmias (complete heart block, ventricular fibrillation, asystole), Outcome 1 Overall (CHB, VF and asystole combined).

Review: Hyperbaric oxygen therapy for acute coronary syndrome

Comparison: 3 Significant dysrhythmias (complete heart block, ventricular fibrillation, asystole)

Outcome: 1 Overall (CHB, VF and asystole combined)




Analysis 3.2. Comparison 3 Significant dysrhythmias (complete heart block, ventricular fibrillation, asystole), Outcome 2 Significant dysrhythmias (complete heart block, ventricular fibrillation or asystole).

Review: Hyperbaric oxygen therapy for acute coronary syndrome

Comparison: 3 Significant dysrhythmias (complete heart block, ventricular fibrillation, asystole)

Outcome: 2 Significant dysrhythmias (complete heart block, ventricular fibrillation or asystole)




Analysis 3.3. Comparison 3 Significant dysrhythmias (complete heart block, ventricular fibrillation, asystole), Outcome 3 Overall best case.

Review: Hyperbaric oxygen therapy for acute coronary syndrome

Comparison: 3 Significant dysrhythmias (complete heart block, ventricular fibrillation, asystole)

Outcome: 3 Overall best case




Analysis 3.4. Comparison 3 Significant dysrhythmias (complete heart block, ventricular fibrillation, asystole), Outcome 4 Overall worst case.

Review: Hyperbaric oxygen therapy for acute coronary syndrome

Comparison: 3 Significant dysrhythmias (complete heart block, ventricular fibrillation, asystole)

Outcome: 4 Overall worst case




Analysis 4.1. Comparison 4 Time to pain relief, Outcome 1 Time to relief of pain.

Review: Hyperbaric oxygen therapy for acute coronary syndrome

Comparison: 4 Time to pain relief

Outcome: 1 Time to relief of pain




Analysis 5.1. Comparison 5 Magnitude of cardiac enzyme changes, Outcome 1 12 hour Plasma Creatine Phosphokinase.

Review: Hyperbaric oxygen therapy for acute coronary syndrome

Comparison: 5 Magnitude of cardiac enzyme changes

Outcome: 1 12 hour Plasma Creatine Phosphokinase




Analysis 5.2. Comparison 5 Magnitude of cardiac enzyme changes, Outcome 2 24 hour Plasma Creatine Phosphate.

Review: Hyperbaric oxygen therapy for acute coronary syndrome

Comparison: 5 Magnitude of cardiac enzyme changes

Outcome: 2 24 hour Plasma Creatine Phosphate




Analysis 5.3. Comparison 5 Magnitude of cardiac enzyme changes, Outcome 3 Maximum Plasma Creatine Phosphate.

Review: Hyperbaric oxygen therapy for acute coronary syndrome

Comparison: 5 Magnitude of cardiac enzyme changes

Outcome: 3 Maximum Plasma Creatine Phosphate




Analysis 6.1. Comparison 6 Improvement in left ventricular function, Outcome 1 Improved contraction in at least one segment (post-HBOT echo).

Review: Hyperbaric oxygen therapy for acute coronary syndrome

Comparison: 6 Improvement in left ventricular function

Outcome: 1 Improved contraction in at least one segment (post-HBOT echo)




Analysis 6.2. Comparison 6 Improvement in left ventricular function, Outcome 2 Left Ventricular Ejection Fraction- % (final estimate).

Review: Hyperbaric oxygen therapy for acute coronary syndrome

Comparison: 6 Improvement in left ventricular function

Outcome: 2 Left Ventricular Ejection Fraction- % (final estimate)




Analysis 7.1. Comparison 7 Length of Stay, Outcome 1 Overall length of stay (days).

Review: Hyperbaric oxygen therapy for acute coronary syndrome

Comparison: 7 Length of Stay

Outcome: 1 Overall length of stay (days)




Analysis 8.1. Comparison 8 Adverse events of therapy, Outcome 1 Total adverse events.


Review: Hyperbaric oxygen therapy for acute coronary syndrome

Comparison: 8 Adverse events of therapy

Outcome: 1 Total adverse events



(Continued ...)



APPENDICES

Appendix I. Search strategies 2004

COCHRANE LIBRARY

- #1 MeSH descriptor HYPERBARIC OXYGENATION explode all trees
- #2 (hyperbaric near/6 oxygen*)
- #3 hbot
- #4 high next pressure next oxygen*
- #5 MeSH descriptor atmosphere exposure chambers this term only
- #6 (monoplace near/6 chamber*)
- #7 (multiplace near/6 chamber*)
- #8 (atmosphere* near/6 chamber*)
- #9 (#1 or #2 or #3 or #4 or #6 or #7 or #8)
- #10 MeSH descriptor MYOCARDIAL ISCHEMIA explode all trees
- #11 myocardial next infarct*
- #12 heart next infarct* 913
- #13 cardiac next infarct* 51
- #14 coronary next thrombosis 212
- #15 acute next coronary 1019
- #16 myocardial next ischaemi* 413
- #17 myocardial next ischemi* 2217
- #18 coronary next disease 6128
- #19 (coronary near/6 disease)
- #20 heart next disease*

#21 unstable next angina
 #22 coronary next arteriosclerosis
 #23 coronary
 #24 ami
 #25 chd
 #26 (ischaemic near/6 heart)
 #27 (ischaemic near/6 heart)
 #28 (#10 or #11 or #12 or #13 or #14 or #15 or #16 or #17 or #18 or #19)
 #29 (#20 or #21 or #22 or #23 or #24 or #25 or #26 or #27)
 #30 (#28 or #29)
 #27 (#30 and #9)

MEDLINE

1 Hyperbaric Oxygenation/
 2 Atmosphere Exposure Chambers/
 3 hyperbaric oxygen\$.tw.
 4 high pressure oxygen.tw.
 5 hbot.tw.
 6 ((monoplace or multiplace) adj5 chamber\$).tw.
 7 or/1-6
 8 exp Myocardial Ischemia/
 9 myocardial infarct\$.tw.
 10 heart infarct\$.tw.
 11 coronary thrombosis.tw.
 12 acute coronary.tw.
 13 myocardial ischaemi\$.tw.
 14 myocardial ischemi\$.tw.
 15 heart disease.tw.
 16 (coronary adj3 disease).tw.
 17 unstable angina.tw.
 18 coronary arteriosclerosis.tw.
 19 (ischaemic adj3 heart).tw.
 20 (ischaemic adj3 heart).tw.
 21 ami.tw.
 22 chd.tw.
 23 or/8-22
 24 7 and 23
 25 randomized controlled trial.pt.
 26 controlled clinical trial.pt.
 27 Randomized controlled trials/
 28 random allocation/
 29 double blind method/
 30 single-blind method/
 31 or/25-30
 32 exp animal/ not humans/
 33 31 not 32
 34 clinical trial.pt.
 35 exp Clinical trials/
 36 (clin\$ adj25 trial\$).ti,ab.
 37 ((singl\$ or doubl\$ or trebl\$ or tripl\$) adj (blind\$ or mask\$)).ti,ab.
 38 placebos/
 39 placebo\$.ti,ab.

40 random\$.ti,ab.
 41 research design/
 42 or/34-41
 43 42 not 32
 44 43 not 33
 45 comparative study.pt.
 46 exp evaluation studies/
 47 follow up studies/
 48 prospective studies/
 49 (control\$ or prospectiv\$ or volunteer\$).ti,ab.
 50 or/45-49
 51 50 not 32
 52 51 not (33 or 44)
 53 33 or 44 or 52
 54 24 and 53

EMBASE

1 Ischemic Heart Disease/
 2 Hyperbaric Oxygen/
 3 hyperbaric oxygen\$.tw.
 4 high pressure oxygen.tw.
 5 hbot.tw.
 6 ((monoplace or multiplace) adj5 chamber\$).tw.
 7 or/2-6
 8 exp Ischemic Heart Disease/
 9 myocardial infarct\$.tw.
 10 heart infarct\$.tw.
 11 coronary thrombosis.tw.
 12 acute coronary.tw.
 13 myocardial ischaemi\$.tw.
 14 myocardial ischemi\$.tw.
 15 heart disease.tw.
 16 (coronary adj3 disease).tw.
 17 unstable angina.tw.
 18 coronary arteriosclerosis.tw.
 19 (ischaemic adj3 heart).tw.
 20 (ischaemic adj3 heart).tw.
 21 ami.tw.
 22 chd.tw.
 23 or/8-22
 24 7 and 23
 25 clinical trial/
 26 random\$.tw.
 27 randomized controlled trial/
 28 trial\$.tw.
 29 follow-up.tw.
 30 double blind procedure/
 31 placebo\$.tw.
 32 placebo/
 33 factorial\$.ti,ab.
 34 (crossover\$ or cross-over\$).ti,ab.
 35 (double\$ adj blind\$).ti,ab.

36 (singl\$ adj blind\$).ti,ab.
 37 assign\$.ti,ab.
 38 allocat\$.ti,ab.
 39 volunteer\$.ti,ab.
 40 Crossover Procedure/
 41 Single Blind Procedure/
 42 or/25-41
 43 exp animal/
 44 nonhuman/
 45 exp animal experiment/
 46 or/43-45
 47 exp human/
 48 46 not 47
 49 42 not 48
 50 24 and 49

CINAHL

1 Hyperbaric Oxygenation/
 2 exp Myocardial Ischemia/
 3 Hyperbaric Oxygenation/
 4 hyperbaric oxygen\$.tw.
 5 high pressure oxygen.tw.
 6 hbot.tw.
 7 ((monoplace or multiplace) adj5 chamber\$).tw.
 8 or/3-7
 9 exp Myocardial Ischemia/
 10 myocardial infarct\$.tw.
 11 heart infarct\$.tw.
 12 coronary thrombosis.tw.
 13 acute coronary.tw.
 14 myocardial ischaemi\$.tw.
 15 myocardial ischemi\$.tw.
 16 heart disease.tw.
 17 (coronary adj3 disease).tw.
 18 unstable angina.tw.
 19 coronary arteriosclerosis.tw.
 20 (ischaemic adj3 heart).tw.
 21 (ischaemic adj3 heart).tw.
 22 ami.tw.
 23 chd.tw.
 24 or/9-23
 25 8 and 24

Appendix 2. Search strategies 2007

COCHRANE LIBRARY

#1 MeSH descriptor HYPERBARIC OXYGENATION explode all trees
#2 (hyperbaric near/6 oxygen*)
#3 hbot
#4 high next pressure next oxygen*
#5 MeSH descriptor atmosphere exposure chambers this term only
#6 (monoplace near/6 chamber*)
#7 (multiplace near/6 chamber*)
#8 (atmosphere* near/6 chamber*)
#9 (#1 or #2 or #3 or #4 or #6 or #7 or #8)
#10 MeSH descriptor MYOCARDIAL ISCHEMIA explode all trees
#11 myocardial next infarct*
#12 heart next infarct* 913
#13 cardiac next infarct* 51
#14 coronary next thrombosis 212
#15 acute next coronary 1019
#16 myocardial next ischaemi* 413
#17 myocardial next ischemi* 2217
#18 coronary next disease 6128
#19 (coronary near/6 disease)
#20 heart next disease*
#21 unstable next angina
#22 coronary next arteriosclerosis
#23 coronary
#24 ami
#25 chd
#26 (ischaemic near/6 heart)
#27 (ischaemic near/6 heart)
#28 (#10 or #11 or #12 or #13 or #14 or #15 or #16 or #17 or #18 or #19)
#29 (#20 or #21 or #22 or #23 or #24 or #25 or #26 or #27)
#30 (#28 or #29)
#27 (#30 and #9)

MEDLINE

1 Hyperbaric Oxygenation/
2 Atmosphere Exposure Chambers/
3 hyperbaric oxygen\$.tw.
4 high pressure oxygen.tw.
5 hbot.tw.
6 ((monoplace or multiplace) adj5 chamber\$).tw.
7 or/1-6
8 exp Myocardial Ischemia/
9 myocardial infarct\$.tw.
10 heart infarct\$.tw.
11 coronary thrombosis.tw.
12 acute coronary.tw.
13 myocardial ischaemi\$.tw.
14 myocardial ischemi\$.tw.
15 heart disease.tw.

16 (coronary adj3 disease).tw.
 17 unstable angina.tw.
 18 coronary arteriosclerosis.tw.
 19 (ischaemic adj3 heart).tw.
 20 (ischaemic adj3 heart).tw.
 21 ami.tw.
 22 chd.tw.
 23 or/8-22
 24 7 and 23
 25 randomized controlled trial.pt.
 26 controlled clinical trial.pt.
 27 Randomized controlled trials/
 28 random allocation/
 29 double blind method/
 30 single-blind method/
 31 or/25-30
 32 exp animal/ not humans/
 33 31 not 32
 34 clinical trial.pt.
 35 exp Clinical trials/
 36 (clin\$ adj25 trial\$).ti,ab.
 37 ((singl\$ or doubl\$ or trebl\$ or tripl\$) adj (blind\$ or mask\$)).ti,ab.
 38 placebos/
 39 placebo\$.ti,ab.
 40 random\$.ti,ab.
 41 research design/
 42 or/34-41
 43 42 not 32
 44 43 not 33
 45 comparative study.pt.
 46 exp evaluation studies/
 47 follow up studies/
 48 prospective studies/
 49 (control\$ or prospectiv\$ or volunteer\$).ti,ab.
 50 or/45-49
 51 50 not 32
 52 51 not (33 or 44)
 53 33 or 44 or 52
 54 24 and 53
 55 limit 54 to yr="2004 - 2007"

EMBASE

1 Ischemic Heart Disease/
 2 Hyperbaric Oxygen/
 3 hyperbaric oxygen\$.tw.
 4 high pressure oxygen.tw.
 5 hbot.tw.
 6 ((monoplace or multiplace) adj5 chamber\$).tw.
 7 or/2-6
 8 exp Ischemic Heart Disease/
 9 myocardial infarct\$.tw.
 10 heart infarct\$.tw.

11 coronary thrombosis.tw.
 12 acute coronary.tw.
 13 myocardial ischaemi\$.tw.
 14 myocardial ischemi\$.tw.
 15 heart disease.tw.
 16 (coronary adj3 disease).tw.
 17 unstable angina.tw.
 18 coronary arteriosclerosis.tw.
 19 (ischaemic adj3 heart).tw.
 20 (ischaemic adj3 heart).tw.
 21 ami.tw.
 22 chd.tw.
 23 or/8-22
 24 7 and 23
 25 clinical trial/
 26 random\$.tw.
 27 randomized controlled trial/
 28 trial\$.tw.
 29 follow-up.tw.
 30 double blind procedure/
 31 placebo\$.tw.
 32 placebo/
 33 factorial\$.ti,ab.
 34 (crossover\$ or cross-over\$).ti,ab.
 35 (double\$ adj blind\$).ti,ab.
 36 (singl\$ adj blind\$).ti,ab.
 37 assign\$.ti,ab.
 38 allocat\$.ti,ab.
 39 volunteer\$.ti,ab.
 40 Crossover Procedure/
 41 Single Blind Procedure/
 42 or/25-41
 43 exp animal/
 44 nonhuman/
 45 exp animal experiment/
 46 or/43-45
 47 exp human/
 48 46 not 47
 49 42 not 48
 50 24 and 49
 51 limit 50 to yr="2004 - 2007"

CINAHL

1 Hyperbaric Oxygenation/
 2 exp Myocardial Ischemia/
 3 Hyperbaric Oxygenation/
 4 hyperbaric oxygen\$.tw.
 5 high pressure oxygen.tw.
 6 hbot.tw.
 7 ((monoplace or multiplace) adj5 chamber\$).tw.
 8 or/3-7
 9 exp Myocardial Ischemia/

10 myocardial infarct\$.tw.
 11 heart infarct\$.tw.
 12 coronary thrombosis.tw.
 13 acute coronary.tw.
 14 myocardial ischaemi\$.tw.
 15 myocardial ischemi\$.tw.
 16 heart disease.tw.
 17 (coronary adj3 disease).tw.
 18 unstable angina.tw.
 19 coronary arteriosclerosis.tw.
 20 (ischaemic adj3 heart).tw.
 21 (ischaemic adj3 heart).tw.
 22 ami.tw.
 23 chd.tw.
 24 or/9-23
 25 8 and 24
 26 limit 25 to yr="2007 - 2014"

Appendix 3. Search strategies 2010

CENTRAL

#1 MeSH descriptor HYPERBARIC OXYGENATION explode all trees
 #2 (hyperbaric near/6 oxygen*)
 #3 hbot
 #4 high next pressure next oxygen*
 #5 MeSH descriptor atmosphere exposure chambers this term only
 #6 (monoplace near/6 chamber*)
 #7 (multiplace near/6 chamber*)
 #8 (atmosphere* near/6 chamber*)
 #9 (#1 or #2 or #3 or #4 or #6 or #7 or #8)
 #10 MeSH descriptor MYOCARDIAL ISCHEMIA explode all trees
 #11 myocardial next infarct*
 #12 heart next infarct*
 #13 cardiac next infarct*
 #14 coronary next thrombosis
 #15 acute next coronary
 #16 myocardial next ischaemi*
 #17 myocardial next ischemi*
 #18 coronary next disease
 #19 (coronary near/6 disease)
 #20 heart next disease*
 #21 unstable next angina
 #22 coronary next arteriosclerosis
 #23 coronary
 #24 ami
 #25 chd
 #26 (ischaemic near/6 heart)
 #27 (ischaemic near/6 heart)
 #28 (#10 or #11 or #12 or #13 or #14 or #15 or #16 or #17 or #18 or #19)
 #29 (#20 or #21 or #22 or #23 or #24 or #25 or #26 or #27)

#30 (#28 or #29)

#27 (#30 and #9)

OID MEDLINE

1. Hyperbaric Oxygenation/
2. Atmosphere Exposure Chambers/
3. hyperbaric oxygen\$.tw.
4. high pressure oxygen.tw.
5. hbot.tw.
6. ((monoplace or multiplace) adj5 chamber\$).tw.
7. or/1-6
8. exp Myocardial Ischemia/
9. myocardial infarct\$.tw.
10. heart infarct\$.tw.
11. coronary thrombosis.tw.
12. acute coronary.tw.
13. myocardial ischaemi\$.tw.
14. myocardial ischemi\$.tw.
15. heart disease.tw.
16. (coronary adj3 disease).tw.
17. unstable angina.tw.
18. coronary arteriosclerosis.tw.
19. (ischaemic adj3 heart).tw.
20. (ischaemic adj3 heart).tw.
21. ami.tw.
22. chd.tw.
23. or/8-22
24. 7 and 23
25. randomized controlled trial.pt.
26. controlled clinical trial.pt.
27. Randomized controlled trials/
28. Random Allocation/
29. Double-Blind Method/
30. single-blind method/
31. or/25-30
32. exp animal/ not humans/
33. 31 not 32
34. clinical trial.pt.
35. exp Clinical Trials as Topic/
36. (clin\$ adj25 trial\$).ti,ab.
37. ((singl\$ or doubl\$ or trebl\$ or tripl\$) adj (blind\$ or mask\$)).ti,ab.
38. Placebos/
39. placebo\$.ti,ab.
40. random\$.ti,ab.
41. Research Design/
42. or/34-41
43. 42 not 32
44. 43 not 33
45. comparative study.pt.
46. evaluation studies/
47. Follow-Up Studies/
48. Prospective Studies/

49. (control\$ or prospectiv\$ or volunteer\$).ti,ab.
50. or/45-49
51. 50 not 32
52. 51 not (33 or 44)
53. 33 or 44 or 52
54. 53 and 24
55. (200909\$ or 200910\$ or 200911\$ or 200912\$ or 2010\$).ed.
56. 54 and 55

OVID EMBASE

- 1 Hyperbaric Oxygen/
- 2 hyperbaric oxygen\$.tw.
- 3 high pressure oxygen.tw.
- 4 hbot.tw.
- 5 ((monoplace or multiplace) adj5 chamber\$).tw.
- 6 or/1-5
- 7 exp Ischemic Heart Disease/
- 8 myocardial infarct\$.tw.
- 9 heart infarct\$.tw.
- 10 coronary thrombosis.tw.
- 11 acute coronary.tw.
- 12 myocardial ischaemi\$.tw.
- 13 myocardial ischemi\$.tw.
- 14 heart disease.tw.
- 15 (coronary adj3 disease).tw.
- 16 unstable angina.tw.
- 17 coronary arteriosclerosis.tw.
- 18 (ischaemic adj3 heart).tw.
- 19 (ischaemic adj3 heart).tw.
- 20 ami.tw.
- 21 chd.tw.
- 22 or/7-21
- 23 6 and 22
- 24 clinical trial/
- 25 random\$.tw.
- 26 randomized controlled trial/
- 27 trial\$.tw.
- 28 follow-up.tw.
- 29 double blind procedure/
- 30 placebo\$.tw.
- 31 placebo/
- 32 factorial\$.ti,ab.
- 33 (crossover\$ or cross-over\$).ti,ab.
- 34 (double\$ adj blind\$).ti,ab.
- 35 (singl\$ adj blind\$).ti,ab.
- 36 assign\$.ti,ab.
- 37 allocat\$.ti,ab.
- 38 volunteer\$.ti,ab.
- 39 Crossover Procedure/
- 40 Single Blind Procedure/
- 41 or/24-40
- 42 exp animal/ not exp human/

43 41 not 42
 44 23 and 43
 45. ("200938" or "200939" or 20094\$ or 20095\$ or "2010").em.
 46. 44 and 45

EBSCO CINAHL

S22 S15 and S21
 S21 S16 or S17 or S18 or S19 or S20
 S20 EM 200912
 S19 EM 200911
 S18 EM 200910
 S17 EM 200909
 S16 EM 2010
 S15 S7 and S14
 S14 S8 or S9 or S10 or S11 or S12 or S13
 S13 monoplace chamber
 S12 multiplace chamber
 S11 high pressure oxygen
 S10 HBOT
 S9 hyperbaric oxygen*
 S8 MH "Hyperbaric Oxygenation"
 S7 S1 or S2 or S3 or S4 or S5 or S6
 S6 "heart infarct*"
 S5 "unstable angina"
 S4 coronary
 S3 ami
 S2 myocardial
 S1 MH "myocardial ischemia+"

Appendix 4. Search strategies 2014

CENTRAL

#1 MeSH descriptor HYPERBARIC OXYGENATION explode all trees
 #2 (hyperbaric near/6 oxygen*)
 #3 hbot
 #4 high next pressure next oxygen*
 #5 MeSH descriptor atmosphere exposure chambers this term only
 #6 (monoplace near/6 chamber*)
 #7 (multiplace near/6 chamber*)
 #8 (atmosphere* near/6 chamber*)
 #9 (#1 or #2 or #3 or #4 or #6 or #7 or #8)
 #10 MeSH descriptor MYOCARDIAL ISCHEMIA explode all trees
 #11 myocardial next infarct*
 #12 heart next infarct*
 #13 cardiac next infarct*
 #14 coronary next thrombosis
 #15 acute next coronary
 #16 myocardial next ischaemi*
 #17 myocardial next ischemi*

#18 coronary next disease
 #19 (coronary near/6 disease)
 #20 heart next disease*
 #21 unstable next angina
 #22 coronary next arteriosclerosis
 #23 coronary
 #24 ami
 #25 chd
 #26 (ischaemic near/6 heart)
 #27 (ischaemic near/6 heart)
 #28 (#10 or #11 or #12 or #13 or #14 or #15 or #16 or #17 or #18 or #19)
 #29 (#20 or #21 or #22 or #23 or #24 or #25 or #26 or #27)
 #30 (#28 or #29)
 #27 (#30 and #9)

OVID MEDLINE

1. Hyperbaric Oxygenation/
 2. Atmosphere Exposure Chambers/
 3. hyperbaric oxygen\$.tw.
 4. high pressure oxygen.tw.
 5. hbot.tw.
 6. ((monoplace or multiplace) adj5 chamber\$).tw.
 7. or/1-6
 8. exp Myocardial Ischemia/
 9. myocardial infarct\$.tw.
 10. heart infarct\$.tw.
 11. coronary thrombosis.tw.
 12. acute coronary.tw.
 13. myocardial ischaemi\$.tw.
 14. myocardial ischemi\$.tw.
 15. heart disease.tw.
 16. (coronary adj3 disease).tw.
 17. unstable angina.tw.
 18. coronary arteriosclerosis.tw.
 19. (ischaemic adj3 heart).tw.
 20. (ischaemic adj3 heart).tw.
 21. ami.tw.
 22. chd.tw.
 23. or/8-22
 24. 7 and 23
 25. randomized controlled trial.pt.
 26. controlled clinical trial.pt.
 27. Randomized controlled trials/
 28. Random Allocation/
 29. Double-Blind Method/
 30. single-blind method/
 31. or/25-30
 32. exp animal/ not humans/
 33. 31 not 32
 34. clinical trial.pt.
 35. exp Clinical Trials as Topic/
 36. (clin\$ adj25 trial\$).ti,ab.

37. ((singl\$ or doubl\$ or trebl\$ or tripl\$) adj (blind\$ or mask\$)).ti,ab.
 38. Placebos/
 39. placebo\$.ti,ab.
 40. random\$.ti,ab.
 41. Research Design/
 42. or/34-41
 43. 42 not 32
 44. 43 not 33
 45. comparative study.pt.
 46. evaluation studies/
 47. Follow-Up Studies/
 48. Prospective Studies/
 49. (control\$ or prospectiv\$ or volunteer\$).ti,ab.
 50. or/45-49
 51. 50 not 32
 52. 51 not (33 or 44)
 53. 33 or 44 or 52
 54. 53 and 24
 55. (2010\$ or 2011\$ or 2012\$ or 2013\$ or 2014\$).ed.
 56. 54 and 55

OVID EMBASE

1 Hyperbaric Oxygen/
 2 hyperbaric oxygen\$.tw.
 3 high pressure oxygen.tw.
 4 hbot.tw.
 5 ((monoplace or multiplace) adj5 chamber\$).tw.
 6 or/1-5
 7 exp Ischemic Heart Disease/
 8 myocardial infarct\$.tw.
 9 heart infarct\$.tw.
 10 coronary thrombosis.tw.
 11 acute coronary.tw.
 12 myocardial ischaemi\$.tw.
 13 myocardial ischemi\$.tw.
 14 heart disease.tw.
 15 (coronary adj3 disease).tw.
 16 unstable angina.tw.
 17 coronary arteriosclerosis.tw.
 18 (ischaemic adj3 heart).tw.
 19 (ischaemic adj3 heart).tw.
 20 ami.tw.
 21 chd.tw.
 22 or/7-21
 23 6 and 22
 24 clinical trial/
 25 random\$.tw.
 26 randomized controlled trial/
 27 trial\$.tw.
 28 follow-up.tw.
 29 double blind procedure/
 30 placebo\$.tw.

31 placebo/
 32 factorial\$.ti,ab.
 33 (crossover\$ or cross-over\$).ti,ab.
 34 (double\$ adj blind\$).ti,ab.
 35 (singl\$ adj blind\$).ti,ab.
 36 assign\$.ti,ab.
 37 allocat\$.ti,ab.
 38 volunteer\$.ti,ab.
 39 Crossover Procedure/
 40 Single Blind Procedure/
 41 or/24-40
 42 exp animal/ not exp human/
 43 41 not 42
 44 23 and 43
 45. ("2010" or "2011" or "2012" or "2013" or "2014").em.
 46. 44 and 45

EBSCO CINAHL

S22 S15 and S21
 S21 S16 or S17 or S18 or S19 or S20
 S20 EM 200912
 S19 EM 200911
 S18 EM 200910
 S17 EM 200909
 S16 EM 2010
 S15 S7 and S14
 S14 S8 or S9 or S10 or S11 or S12 or S13
 S13 monoplace chamber
 S12 multiplace chamber
 S11 high pressure oxygen
 S10 HBOT
 S9 hyperbaric oxygen*
 S8 MH "Hyperbaric Oxygenation"
 S7 S1 or S2 or S3 or S4 or S5 or S6
 S6 "heart infarct*"
 S5 "unstable angina"
 S4 coronary
 S3 ami
 S2 myocardial
 S1 MH "myocardial ischemia+"

WHAT'S NEW

Last assessed as up-to-date: 17 September 2014.

Date	Event	Description
8 October 2014	New citation required but conclusions have not changed	No new studies were found for inclusion
14 September 2014	New search has been performed	Searches were re-run on the 14th September 2014. No new citations We have modified both the background and discussion under the suggested subheadings in these sections. The included studies, included data and meta-analyses are unchanged

HISTORY

Protocol first published: Issue 3, 2004

Review first published: Issue 2, 2005

Date	Event	Description
27 August 2010	New search has been performed	The search has been re-run to June 2010. We identified and included one new trial from the updated search
27 August 2010	New citation required and conclusions have changed	A total of six trials are included in this update. There is now some evidence that hyperbaric oxygen therapy reduces the risk of death from acute coronary syndrome
9 September 2008	Amended	Converted to new review format.
6 July 2007	New citation required and conclusions have changed	Substantive amendment

CONTRIBUTIONS OF AUTHORS

Dr. Michael Bennett: Conception, principal author, search strategy, identification of trials, critical appraisal and data extraction. Content expert on hyperbaric medicine and clinical epidemiology. Guarantor of this review.

Dr. Nigel Jepson: Co-author, critical appraisal and data extraction. Content expert on acute coronary syndrome.

Dr. Jan Lehm: Co-author, data extraction. Content expert on hyperbaric medicine.

DECLARATIONS OF INTEREST

None known.

SOURCES OF SUPPORT

Internal sources

- No internal source of support, Australia.

External sources

- No external source of support, Australia.

INDEX TERMS

Medical Subject Headings (MeSH)

Acute Coronary Syndrome [mortality; *therapy]; Angina, Unstable [mortality; *therapy]; Hyperbaric Oxygenation [*mortality]; Myocardial Infarction [mortality; *therapy]; Randomized Controlled Trials as Topic

MeSH check words

Humans