

FabGennix International, Inc.
 9191 Kyser Way Bldg. 4 Suite 400
 Frisco, TX 75033
 Tel: (214)-387-8105, 1-800-786-1236 Fax: (214)-387-0870 Email: info@fabgennix.com Web: www.fabgennix.com

MUC1 Antibody

Catalog Number: PC-MUC

General Product Information

Product	MUC1
Description	Affinity Purified "ADMCKD antibody, ADMCKD1 antibody, Breast carcinoma associated antigen DF3 antibody, Breast carcinoma-associated antigen DF3 antibody, CA 15 3 antibody, CA 15-3 antibody, CA15 3 antibody, CA15 3 antigen antibody, CA15.3 antibody, Cancer antigen 15-3 antibody, Carcinoma associated mucin antibody, Carcinoma-associated mucin antibody, CD 227 antibody, CD227 antibody, CD227 antigen antibody, DF3 antigen antibody, EMA antibody, Episialin antibody, Epithelial basement membrane antigen antibody, Epithelial membrane antigen antibody, Epithelial mucin tandem repeat sequence antibody, H23 antigen antibody, H23AG antibody, HGNC:7508 antibody, KL 6 antibody, KL-6 antibody, KL6 antibody, Krebs von den Lungen-6 antibody, MAM 6 antibody, MAM6 antibody, MCD antibody, MCKD antibody, MCKD1 antibody, Medullary cystic kidney disease 1 (autosomal dominant) antibody, MUC 1 antibody, MUC 1 antibody, MUC-1 antibody, MUC-1/SEC antibody, MUC-1/X antibody, MUC1 antibody, MUC1-alpha antibody, MUC1-beta antibody, MUC1-CT antibody, MUC1-NT antibody, MUC1/ZD antibody, MUC1_HUMAN antibody, Mucin 1 antibody, Mucin 1 precursor antibody, Mucin 1 transmembrane antibody, Mucin 1, cell surface associated antibody, Mucin-1 subunit beta antibody, Mucin1 antibody, Peanut reactive urinary mucin antibody, Peanut-reactive urinary mucin antibody, PEM antibody, PEMT antibody, Polymorphic epithelial mucin antibody, PUM antibody, Tumor associated epithelial membrane antigen antibody, Tumor associated epithelial mucin antibody, Tumor associated mucin antibody, Tumor-associated epithelial membrane antigen antibody, Tumor-associated mucin antibody," Antibody
Verified Applications	WB

Physical Properties

Quantity	100 µg
Volume	Inquire
Form	Western Blot Positive Control
Immunoglobulin	IgG
Concentration	The concentration is set so that 20 µl per lane in Western and blotted with corresponding affinity purified antibody will give a clean band with a high signal to noise ration on the gel.
Purification Method	Affinity Purified
Clonality	Polyclonal
Storage	-20°C for long term storage.

Application Protocol

Western Blot	1:500–1:2,000
--------------	---------------

This product is for research use only. For more information about this product please visit our website at fabgennix.com or contact us today.