

FabGennix International, Inc.
 9191 Kyser Way Bldg. 4 Suite 400
 Frisco, TX 75033
 Tel: (214)-387-8105, 1-800-786-1236 Fax: (214)-387-0870 Email: info@fabgennix.com Web: www.fabgennix.com

Phospho-ErbB 2 Antibody

Catalog Number: PC-PHER2

General Product Information

Product	Phospho-ErbB 2
Description	Affinity Purified "Verb b2 erythroblastic leukemia viral oncogene homolog 2, neuro/glioblastoma derived oncogene homolog antibody, c erb B2 antibody, c erb B2/neu protein antibody, C erbB 2 antibody, C-erbB2 antibody, CD340 antibody, CD340 antigen antibody, Cerb B2/neu protein antibody, CerbB2 antibody, cneu antibody, Erb B2 antibody, erbb2 antibody, ERBB2_HUMAN antibody, Erythroblastosis Virus antibody, HER 2 antibody, HER 2/neu antibody, HER2 antibody, Her2/neu antibody, Herstatin antibody, Metastatic lymph node gene 19 protein antibody, MLN 19 antibody, MLN19 antibody, NEU antibody, NEU proto oncogene antibody, Neuro Glioblastoma Derived Oncogene Homolog antibody, Neuroblastoma/glioblastoma derived oncogene homolog antibody, NGL antibody, p185 ErbB2 antibody, p185erbB2 antibody, Proto-oncogene c-ErbB-2 antibody, Proto-oncogene Neu antibody, Receptor Protein Tyrosine Kinase ErbB2 Precursor antibody, Receptor tyrosine protein kinase erbB 2 antibody, Receptor tyrosine-protein kinase erbB-2 antibody, TKR1 antibody, Tyrosine kinase type cell surface receptor HER2 antibody, Tyrosine kinase-type cell surface receptor HER2 antibody, V erb b2 avian erythroblastic leukemia viral oncogene homolog 2 (neuro/glioblastoma derived oncogene homolog) antibody, v erb b2 erythroblastic leukemia viral oncogene homolog 2 neuro/glioblastoma derived oncogene homolog (avian) antibody, V erb b2 erythroblastic leukemia viral oncogene homolog 2, neuro/glioblastoma derived oncogene homolog (avian) antibody, Verb b2 erythroblastic leukemia viral oncogene homolog 2, neuro/glioblastoma derived oncogene homolog (avian) antibody, verbb2 antibody, " Antibody
Verified Applications	WB

Physical Properties

Quantity	100 µg
Volume	Inquire
Form	Western Blot Positive Control
Immunoglobulin	IgG
Concentration	The concentration is set so that 20 µl per lane in Western and blotted with corresponding affinity purified antibody will give a clean band with a high signal to noise ration on the gel.
Purification Method	Affinity Purified
Clonality	Polyclonal
Storage	-20°C for long term storage.

Application Protocol

Western Blot	1:500–1:2,000
--------------	---------------

This product is for research use only. For more information about this product please visit our website at fabgennix.com or contact us today.