

B.Sc. (CBCS) DEGREE EXAMINATION,
NOVEMBER 2023

Second/Fourth Semester

Mathematics – Allied

VECTOR CALCULUS AND FOURIER SERIES

(For those who joined in July 2017-2020)

Time : Three hours

Maximum : 75 marks

PART A — (10 × 1 = 10 marks)

Answer ALL questions.

Choose the correct answers.

1. The value of $\vec{i} \cdot \vec{i} =$ _____.
- (a) $\vec{0}$ (b) 0
(c) 1 (d) \vec{j}
2. A vector \vec{f} is harmonic vector if _____.
- (a) $\nabla \vec{f} = 0$ (b) $\nabla^2 \vec{f} = 0$
(c) $\nabla \times \vec{f} = 0$ (d) $\nabla^2 \times \vec{f} = 0$

3. The value of $\int_0^2 \int_0^5 xy \, dx \, dy$

- (a) $\frac{21}{2}$ (b) $\frac{63}{4}$
(c) $\frac{64}{3}$ (d) $\frac{22}{2}$

4. The value of $\int_0^2 \int_0^2 \int_0^2 xy^2 z \, dz \, dy \, dx$

- (a) 26 (b) 24
(c) 42 (d) 62

5. If $\vec{f} = x^2 \vec{i} - xy \vec{j}$ and C is the straight line joining the points (0, 0) and (1, 1) then $\int_C \vec{f} \cdot d\vec{r}$ is _____.

- (a) 0 (b) 1
(c) x^2 (d) xy

6. If $\vec{A} = \text{curl } \vec{F}$, then $\iint_S \vec{A} \cdot \vec{n} \, dS =$ _____ for any closed S .

- (a) 0 (b) S
(c) V (d) $3V$

Page 2 Code No. : 20073 E

7. The value of $\iiint_V \nabla \phi \, dV$ is

- (a) $\int_S \vec{f} \cdot dS$ (b) $\iint_S \phi \vec{n} \, dS$
(c) $\iiint_V \phi \, dV$ (d) 0

8. If $\text{curl } \vec{F} = 0$, then $\int_C \vec{F} \cdot d\vec{r} =$

- (a) \vec{n} (b) 0
(c) 1 (d) \vec{r}

9. Which one of the following is an even function?

- (a) x (b) $\sin x$
(c) e^x (d) x^2

10. If $f(x)$ is odd, then $\int_{-a}^a f(x) \, dx$

- (a) $2 \int_a^a f(x) \, dx$ (b) $\int_{-a}^a f(x) \, dx$
(c) 0 (d) 1

Page 3 Code No. : 20073 E

PART B — (5 × 5 = 25 marks)

Answer ALL questions, choosing either (a) or (b).

11. (a) Prove that $\nabla f(r) = \left(\frac{f'(r)}{r} \right) \vec{r}$.

Or

(b) Show that $\text{div} \left(\frac{\vec{r}}{r} \right) = \frac{2}{r}$.

12. (a) Evaluate $\int_0^{\pi/2} \int_0^{1-\cos\theta} r \, dr \, d\theta$.

Or

(b) Evaluate $\int_0^2 \int_0^2 \int_0^2 xy^2 z \, dz \, dy \, dx$.

13. (a) Evaluate $\int_C \vec{f} \cdot d\vec{r}$ where

 $\vec{f} = (x^2 + y^2)\vec{i} + (x^2 + y^2)\vec{j}$ and C is the curve $y = x^2$ joining (0, 0) and (1, 1).

Or

(b) Find the common area between $y^2 = 4x$ and $x^2 = 4y$ using Green's theorem.

Page 4 Code No. : 20073 E
[P.T.O.]

14. (a) Show that $\iint_S \vec{f} \cdot \vec{n} d\vec{S} = \iiint_V a^2 dV$ where $\vec{r} = \phi \vec{a}$ and $\vec{a} = \nabla \phi$ and $\nabla^2 \phi = 0$.

Or

- (b) Evaluate using Stoke's theorem $\int_C (yz dx + zx dy + xy dz)$ where C is the curve $x^2 + y^2 = 1, z = y^2$.

15. (a) In $-\pi < x < \pi$, express $\sinh ax$ as in Fourier series of period 2π .

Or

- (b) Explain the Half range sine series.

PART C — (5 × 8 = 40 marks)

Answer ALL questions, choosing either (a) or (b).

16. (a) If $\nabla \phi = 2xyz^3 \vec{i} + x^2 z^3 \vec{j} + 3x^2 yz^2 \vec{k}$, find $\phi(x, y, z)$ if $\phi(1, -2, 2) = 4$.

Or

- (b) If \vec{f} is solenoidal prove that $\text{curl curl curl curl } \vec{f} = \nabla^2 \vec{f}$.

Page 5 Code No. : 20073 E

17. (a) Find the value of $\iint xy dx dy$ taken over the positive quadrant of the ellipse $\frac{x^2}{a^2} + \frac{y^2}{b^2} = 1$.

Or

- (b) Evaluate $\int_0^{\infty} \int_0^{\infty} e^{-(x^2+y^2)} dx dy$.

18. (a) Evaluate $\int_C \vec{f} \cdot d\vec{r}$ where $\vec{f} = (x^2 + y^2) \vec{i} - 2xy \vec{j}$ and the curve C is the rectangle in the $x-y$ plane bounded by $y = 0, y = b, x = 0, x = a$.

Or

- (b) Evaluate $\iiint_S \vec{f} \cdot \vec{n} dS$ where $\vec{f} = (x+y) \vec{i} - 2x \vec{j} + 2yz \vec{k}$ and S is the surface of the plane $2x + y + 2z = 6$ in the first octant.

19. (a) Verify using Gauss divergence theorem for $\vec{f} = (x^2 - yz) \vec{i} + (y^2 - zx) \vec{j} + (z^2 - xy) \vec{k}$ over the rectangular parallopiped $0 \leq x \leq a, 0 \leq y \leq b, 0 \leq z \leq c$.

Or

- (b) Evaluate $\int_C (e^x dx + 2y dy - dz)$ by using stokes theorem where C is the curve $x^2 + y^2 = 4, z = 2$.

Page 6 Code No. : 20073 E

20. (a) Expand in Fourier series of $f(x) = x \sin x$ for $0 < x < 2\pi$.

Or

- (b) Find the half range Fourier cosine series for $f(x) = x$ in $0 < x < \pi$.

Page 7 Code No. : 20073 E