

Application of Matrices and Determinants MCQ

12th Standard

Maths

- 1) If $|\text{adj}(\text{adj } A)| = |A|^9$, then the order of the square matrix A is
(a) 3 (b) 4 (c) 2 (d) 5
- 2) If A is a 3×3 non-singular matrix such that $AA^T = A^T A$ and $B = A^{-1}A^T$, then $BB^T =$
(a) A (b) B (c) I_3 (d) B^T
- 3) If $A = \begin{bmatrix} 3 & 5 \\ 1 & 2 \end{bmatrix}$, $B = \text{adj } A$ and $C = 3A$, then $\frac{|\text{adj } B|}{|C|} =$
(a) $\frac{1}{3}$ (b) $\frac{1}{9}$ (c) $\frac{1}{4}$ (d) 1
- 4) If $A \begin{bmatrix} 1 & -2 \\ 1 & 4 \end{bmatrix} = \begin{bmatrix} 6 & 0 \\ 0 & 6 \end{bmatrix}$, then A =
(a) $\begin{bmatrix} 1 & -2 \\ 1 & 4 \end{bmatrix}$ (b) $\begin{bmatrix} 1 & 2 \\ -1 & 4 \end{bmatrix}$ (c) $\begin{bmatrix} 4 & 2 \\ -1 & 1 \end{bmatrix}$ (d) $\begin{bmatrix} 4 & -1 \\ 2 & 1 \end{bmatrix}$
- 5) If $A = \begin{bmatrix} 7 & 3 \\ 4 & 2 \end{bmatrix}$, then $9I_2 - A =$
(a) A^{-1} (b) $\frac{A^{-1}}{2}$ (c) $3A^{-1}$ (d) $2A^{-1}$
- 6) If $A = \begin{bmatrix} 2 & 0 \\ 1 & 5 \end{bmatrix}$ and $B = \begin{bmatrix} 1 & 4 \\ 2 & 0 \end{bmatrix}$ then $|\text{adj } (AB)| =$
(a) -40 (b) -80 (c) -60 (d) -20
- 7) If $P = \begin{bmatrix} 1 & x & 0 \\ 1 & 3 & 0 \\ 2 & 4 & -2 \end{bmatrix}$ is the adjoint of 3×3 matrix A and $|A| = 4$, then x is
(a) 15 (b) 12 (c) 14 (d) 11

8) If $A = \begin{bmatrix} 3 & 1 & -1 \\ 2 & -2 & 0 \\ 1 & 2 & -1 \end{bmatrix}$ and $A^{-1} = \begin{bmatrix} a_{11} & a_{12} & a_{13} \\ a_{21} & a_{22} & a_{23} \\ a_{31} & a_{32} & a_{33} \end{bmatrix}$ then the value of a_{23} is

- (a) 0 (b) -2 (c) -3 (d) -1

9) If A, B and C are invertible matrices of some order, then which one of the following is not true?

- (a) $\text{adj } A = |A| A^{-1}$ (b) $\text{adj}(AB) = (\text{adj } A)(\text{adj } B)$ (c) $\det A^{-1} = (\det A)^{-1}$
(d) $(ABC)^{-1} = C^{-1}B^{-1}A^{-1}$

10) If $(AB)^{-1} = \begin{bmatrix} 12 & -17 \\ -19 & 27 \end{bmatrix}$ and $A^{-1} = \begin{bmatrix} 1 & -1 \\ -2 & 3 \end{bmatrix}$, then $B^{-1} =$

- (a) $\begin{bmatrix} 2 & -5 \\ -3 & 8 \end{bmatrix}$ (b) $\begin{bmatrix} 8 & 5 \\ 3 & 2 \end{bmatrix}$ (c) $\begin{bmatrix} 3 & 1 \\ 2 & 1 \end{bmatrix}$ (d) $\begin{bmatrix} 8 & -5 \\ -3 & 2 \end{bmatrix}$

11) If $A^T A^{-1}$ is symmetric, then $A^2 =$

- (a) A^{-1} (b) $(A^T)^2$ (c) A^T (d) $(A^{-1})^2$

12) If A is a non-singular matrix such that $A^{-1} = \begin{bmatrix} 5 & 3 \\ -2 & -1 \end{bmatrix}$, then $(A^T)^{-1} =$

- (a) $\begin{bmatrix} -5 & 3 \\ 2 & 1 \end{bmatrix}$ (b) $\begin{bmatrix} 5 & 3 \\ -2 & -1 \end{bmatrix}$ (c) $\begin{bmatrix} -1 & -3 \\ 2 & 5 \end{bmatrix}$ (d) $\begin{bmatrix} 5 & -2 \\ 3 & -1 \end{bmatrix}$

13) If $A = \begin{bmatrix} \frac{3}{5} & \frac{4}{5} \\ x & \frac{3}{5} \end{bmatrix}$ and $A^T = A^{-1}$, then the value of x is

- (a) $\frac{-4}{5}$ (b) $\frac{-3}{5}$ (c) $\frac{3}{5}$ (d) $\frac{4}{5}$

14) If $A = \begin{bmatrix} 1 & \tan \frac{\theta}{2} \\ -\tan \frac{\theta}{2} & 1 \end{bmatrix}$ and $AB = I_2$, then $B =$

- (a) $\left(\cos^2 \frac{\theta}{2}\right) A$ (b) $\left(\cos^2 \frac{\theta}{2}\right) A^T$ (c) $(\cos^2 \theta) I$ (d) $(\sin^2 \frac{\theta}{2}) A$

15) If $A = \begin{bmatrix} \cos \theta & \sin \theta \\ -\sin \theta & \cos \theta \end{bmatrix}$ and $A(\text{adj } A) = \begin{bmatrix} k & 0 \\ 0 & k \end{bmatrix}$, then k =

- (a) 0 (b) $\sin \theta$ (c) $\cos \theta$ (d) 1

16) If $A = \begin{bmatrix} 2 & 3 \\ 5 & -2 \end{bmatrix}$ be such that $\lambda A^{-1} = A$, then λ is

- (a) 17 (b) 14 (c) 19 (d) 21

JOIN 10TH & 12TH PAID WHATSAPP GROUP

DEC 1 2025 TO TILL FINAL 2026 MARCH EXAM
UPLOAD DAILY ONE SUBJECT FULL TEST PAPER

ONE TIME FEES RS.750 ONLY

ALL TEST PAPERS WITH ANSWERS UPLOAD IN YOUR WHATSAPP

FREE USERS CHECK ALL SAME PAPERS UPLOAD IN MY WEBSITES. YOU CAN DOWNLOAD AND PRACTICE

RAVI TEST PAPERS & NOTES
WHATSAPP – 8056206308

CHECK FREE TEST PAPERS www.ravitestpapers.com & www.ravitestpapers.in

JOIN PAID WHATSAPP GROUP WITH PDF ANSWERS
ONE TIME FEES RS.750 TILL FINAL BOARD EXAM

17) If $\text{adj } A = \begin{bmatrix} 2 & 3 \\ 4 & -1 \end{bmatrix}$ and $\text{adj } B = \begin{bmatrix} 1 & -2 \\ -3 & 1 \end{bmatrix}$ then $\text{adj } (AB)$ is

- (a) $\begin{bmatrix} -7 & -1 \\ 7 & -9 \end{bmatrix}$ (b) $\begin{bmatrix} -6 & 5 \\ -2 & -10 \end{bmatrix}$ (c) $\begin{bmatrix} -7 & 7 \\ -1 & -9 \end{bmatrix}$ (d) $\begin{bmatrix} -6 & -2 \\ 5 & -10 \end{bmatrix}$

18) The rank of the matrix $\begin{bmatrix} 1 & 2 & 3 & 4 \\ 2 & 4 & 6 & 8 \\ -1 & -2 & -3 & -4 \end{bmatrix}$ is

- (a) 1 (b) 2 (c) 4 (d) 3

19) If $x^a y^b = e^m$, $x^c y^d = e^n$, $\Delta_1 = \begin{vmatrix} m & b \\ n & d \end{vmatrix}$, $\Delta_2 = \begin{vmatrix} a & m \\ c & n \end{vmatrix}$, $\Delta_3 = \begin{vmatrix} a & b \\ c & d \end{vmatrix}$, then the values of x and y are respectively,

- (a) $e^{(\Delta_2 / \Delta_1)}$, $e^{(\Delta_3 / \Delta_1)}$ (b) $\log (\Delta_1 / \Delta_3)$, $\log (\Delta_2 / \Delta_3)$
 (c) $\log (\Delta_2 / \Delta_1)$, $\log (\Delta_3 / \Delta_1)$ (d) $e^{(\Delta_1 / \Delta_3)}$, $e^{(\Delta_2 / \Delta_3)}$

20) Which of the following is/are correct?

- (i) Adjoint of a symmetric matrix is also a symmetric matrix.
 (ii) Adjoint of a diagonal matrix is also a diagonal matrix.
 (iii) If A is a square matrix of order n and λ is a scalar, then $\text{adj}(\lambda A) = \lambda^n \text{adj}(A)$.
 (iv) $A(\text{adj} A) = (\text{adj} A)A = |A| I$

- (a) Only (i) (b) (ii) and (iii) (c) (iii) and (iv) (d) (i), (ii) and (iv)

21) If $\rho(A) = \rho([A | B])$, then the system $AX = B$ of linear equations is

- (a) consistent and has a unique solution (b) consistent
 (c) consistent and has infinitely many solution (d) inconsistent

22) If $0 \leq \theta \leq \pi$ and the system of equations $x + (\sin\theta)y - (\cos\theta)z = 0$, $(\cos\theta)x - y + z = 0$, $(\sin\theta)x + y - z = 0$ has a non-trivial solution then θ is

- (a) $\frac{2\pi}{3}$ (b) $\frac{3\pi}{4}$ (c) $\frac{5\pi}{6}$ (d) $\frac{\pi}{4}$

23) The augmented matrix of a system of linear equations is

$$\begin{bmatrix} 1 & 2 & 7 & 3 \\ 0 & 1 & 4 & 6 \\ 0 & 0 & \lambda - 7 & \mu + 5 \end{bmatrix}. \text{ The system has infinitely many solutions if}$$

- (a) $\lambda = 7, \mu \neq -5$ (b) $\lambda = -7, \mu = 5$ (c) $\lambda \neq 7, \mu \neq -5$ (d) $\lambda = 7, \mu = -5$

24) Let $A = \begin{bmatrix} 2 & -1 & 1 \\ -1 & 2 & -1 \\ 1 & -1 & 2 \end{bmatrix}$ and $4B = \begin{bmatrix} 3 & 1 & -1 \\ 1 & 3 & x \\ -1 & 1 & 3 \end{bmatrix}$. If B is the inverse of A , then the value of x is

- (a) 2 (b) 4 (c) 3 (d) 1

25) If $A = \begin{bmatrix} 3 & -3 & 4 \\ 2 & -3 & 4 \\ 0 & -1 & 1 \end{bmatrix}$, then $\text{adj}(\text{adj } A)$ is

- (a) $\begin{bmatrix} 3 & -3 & 4 \\ 2 & -3 & 4 \\ 0 & -1 & 1 \end{bmatrix}$ (b) $\begin{bmatrix} 6 & -6 & 8 \\ 4 & -6 & 8 \\ 0 & -2 & 2 \end{bmatrix}$ (c) $\begin{bmatrix} -3 & 3 & -4 \\ -2 & 3 & -4 \\ 0 & 1 & -1 \end{bmatrix}$ (d) $\begin{bmatrix} 3 & -3 & 4 \\ 0 & -1 & 1 \\ 2 & -3 & 4 \end{bmatrix}$

26) The system of linear equations $x + y + z = 6$, $x + 2y + 3z = 14$ and $2x + 5y + \lambda z = \mu$ ($\lambda, \mu \in \mathbb{R}$) is consistent with unique solution if _____

- (a) $\lambda = 8$ (b) $\lambda = 8, \mu \neq 36$ (c) $\lambda \neq 8$ (d) none

27) If the system of equations $x = cy + bz$, $y = az + cx$ and $z = bx + ay$ has a non-trivial solution then _____

- (a) $a^2 + b^2 + c^2 = 1$ (b) $abc \neq 1$ (c) $a + b + c = 0$ (d) $a^2 + b^2 + c^2 + 2abc = 1$

28) Let A be a 3×3 matrix and B its adjoint matrix. If $|B| = 64$, then $|A| =$ _____

- (a) ± 2 (b) ± 4 (c) ± 8 (d) ± 12

29) If A^T is the transpose of a square matrix A , then _____

- (a) $|A| \neq |A^T|$ (b) $|A| = |A^T|$ (c) $|A| + |A^T| = 0$ (d) $|A| = |A^T|$ only

30) The number of solutions of the system of equations $2x + y = 4$, $x - 2y = 2$, $3x + 5y = 6$ is _____

- (a) 0 (b) 1 (c) 2 (d) infinitely many

31) If A is a square matrix that $|A| = 2$, then for any positive integer n , $|A^n| =$ _____

- (a) 0 (b) $2n$ (c) 2^n (d) n^2

32) The system of linear equations $x + y + z = 2$, $2x + y - z = 3$, $3x + 2y + kz =$ has a unique solution if _____

- (a) $k \neq 0$ (b) $-1 < k < 1$ (c) $-2 < k < 2$ (d) $k = 0$

33) If A is a square matrix of order n , then $|\text{adj } A| =$ _____

- (a) $|A|^{n-1}$ (b) $|A|^{n-2}$ (c) $|A|^n$ (d) None

34) If the system of equations $x + 2y - 3z = 2$, $(k + 3)z = 3$, $(2k + 1)y + z = 2$ is inconsistent then k is _____

- (a) $-3, -\frac{1}{2}$ (b) $-\frac{1}{2}$ (c) 1 (d) 2

35) If $A = \begin{pmatrix} \cos x & \sin x \\ -\sin x & \cos x \end{pmatrix}$ and $A(\text{adj } A) = \lambda \begin{pmatrix} 1 & 0 \\ 0 & 1 \end{pmatrix}$ then λ is _____

- (a) $\sin x \cos x$ (b) 1 (c) 2 (d) none

36) If A is a matrix of order $m \times n$, then $\rho(A)$ is _____

- (a) m (b) n (c) $\leq \min(m, n)$ (d) $\geq \min(m, n)$

37) The system of equations $x + 2y + 3z = 1$, $x - y + 4z = 0$, $2x + y + 7z = 1$ has _____

- (a) One solution (b) Two solution (c) No solution
(d) Infinitely many solution

38) If $\rho(A) = \rho([A/B]) = \text{number of unknowns}$, then the system is _____--

- (a) consistent and has infinitely many solutions (b) consistent
(c) inconsistent (d) consistent and has unique solution

39) Which of the following is not an elementary transformation?

- (a) $R_i \leftrightarrow R_j$ (b) $R_i \rightarrow 2R_i + R_j$ (c) $C_j \rightarrow C_j + C_i$ (d) $R_i \rightarrow R_i + C_j$

40) If $\rho(A) = r$ then which of the following is correct?

- (a) all the minors of order n which do not vanish
(b) 'A' has at least one minor of order r which does not vanish and all higher order minors vanish
(c) 'A' has at least one $(r + 1)$ order minor which vanish
(d) all $(r + 1)$ and higher order minors should not vanish

41) Every homogeneous system _____

- (a) Is always consistent (b) Has only trivial solution
(c) Has infinitely many solution (d) Need not be consistent

42) If $\rho(A) \neq \rho([A/B])$, then the system is _____

- (a) consistent and has infinitely many solutions
(b) consistent and has a unique solution (c) consistent (d) inconsistent

43) In the non - homogeneous system of equations with 3 unknowns if $\rho(A) = \rho([A/B]) = 2$, then the system has _____

- (a) unique solution (b) one parameter family of solution
(c) two parameter family of solutions (d) inconsistent

44) Cramer's rule is applicable only when _____

- (a) $\Delta \neq 0$ (b) $\Delta = 0$ (c) $\Delta = 0, \Delta_x = 0$ (d) $\Delta_x = \Delta_y = \Delta_z = 0$

45) In a homogeneous system if $\rho(A) = \rho([A|0]) < \text{the number of unknowns}$ then the system has _____

- (a) trivial solution (b) only non - trivial solution (c) no solution
(d) trivial solution and infinitely many non - trivial solutions

46) In the system of equations with 3 unknowns, if $\Delta = 0$, and one of Δ_x , Δ_y of Δ_z is non zero then the system is _____

- (a) Consistent (b) inconsistent
(c) consistent with one parameter family of solutions
(d) consistent with two parameter family of solutions

47) In the system of liner equations with 3 unknowns If $\rho(A) = \rho([A|B]) = 1$, the system has _____

- (a) unique solution (b) inconsistent
(c) consistent with 2 parameter -family of solution
(d) consistent with one parameter family of solution.

48) If $A = [2 \ 0 \ 1]$ then the rank of AA^T is _____

- (a) 1 (b) 2 (c) 3 (d) 0

49) If A is a non-singular matrix then $|A^{-1}| =$ _____

- (a) $\left| \frac{1}{A^2} \right|$ (b) $\frac{1}{|A^2|}$ (c) $\left| \frac{1}{A} \right|$ (d) $\frac{1}{|A|}$

50) In a square matrix the minor M_{ij} and the co-factor A_{ij} of and element a_{ij} are related by _____

- (a) $A_{ij} = -M_{ij}$ (b) $A_{ij} = M_{ij}$ (c) $A_{ij} = (-1)^{i+j} M_{ij}$ (d) $A_{ij} = (-1)^{i-j} M_{ij}$

51) Let $A = \begin{bmatrix} 4 & 4k & k \\ 0 & k & 4k \\ 0 & 0 & 4 \end{bmatrix}$ If $\det(A^2) = 16$ then $|k|$ is _____

- (a) 1 (b) $\frac{1}{4}$ (c) 4 (d) 4^2

52) Let $1, \omega, \omega^2$ are cube roots of unity then

$$\begin{vmatrix} a & a^2 & a^3 - 1 \\ a^\omega & a^{2\omega} & a^{3\omega} - 1 \\ a^{\omega^2} & a^{2\omega^2} & a^{3\omega^2} - 1 \end{vmatrix} = \underline{\hspace{2cm}}$$

- (a) 0 (b) a (c) a^2 (d) a^3

53) If $P = \begin{bmatrix} \frac{\sqrt{3}}{2} & \frac{1}{2} \\ -\frac{1}{2} & \frac{\sqrt{3}}{2} \end{bmatrix}$, $A = \begin{bmatrix} 1 & 1 \\ 0 & 1 \end{bmatrix}$ and $Q = PAP^T$, then $P^T Q^{2013} P =$

- (a) $\begin{bmatrix} 1 & 2013 \\ 0 & 1 \end{bmatrix}$ (b) $\begin{bmatrix} 4 + 2013\sqrt{3} & 6039 \\ 2012 & 4 - 2013\sqrt{3} \end{bmatrix}$ (c) $\frac{1}{4} \begin{bmatrix} 2 + \sqrt{3} & 1 \\ -1 & 2 - \sqrt{3} \end{bmatrix}$
(d) $\frac{1}{4} \begin{bmatrix} 2012 & 2 - \sqrt{3} \\ 2 + \sqrt{3} & 2012 \end{bmatrix}$

JOIN 10TH & 12TH PAID WHATSAPP GROUP

DEC 1 2025 TO TILL FINAL 2026 MARCH EXAM
UPLOAD DAILY ONE SUBJECT FULL TEST PAPER

ONE TIME FEES RS.750 ONLY

ALL TEST PAPERS WITH ANSWERS UPLOAD IN YOUR WHATSAPP

FREE USERS CHECK ALL SAME PAPERS UPLOAD IN MY WEBSITES. YOU CAN DOWNLOAD AND PRACTICE

RAVI TEST PAPERS & NOTES

WHATSAPP – 8056206308

CHECK FREE TEST PAPERS www.ravitestpapers.com & www.ravitestpapers.in

JOIN 10TH & 12TH PAID WHATSAPP GROUP

DEC 1 2025 TO TILL FINAL 2026 MARCH EXAM
UPLOAD DAILY ONE SUBJECT FULL TEST PAPER

ONE TIME FEES RS.750 ONLY

ALL TEST PAPERS WITH ANSWERS UPLOAD IN YOUR WHATSAPP

FREE USERS CHECK ALL SAME PAPERS UPLOAD IN MY WEBSITES. YOU CAN DOWNLOAD AND PRACTICE

RAVI TEST PAPERS & NOTES

WHATSAPP – 8056206308

CHECK FREE TEST PAPERS www.ravitestpapers.com & www.ravitestpapers.in

54) If $\bar{A} = \begin{bmatrix} -1 & 2-3i & 3+4i \\ 2+3i & 5 & 1+i \\ 3-4i & 1-i & 4 \end{bmatrix}$ then $\det A$ is _____

(a) purely real (b) purely imaginary (c) complex number (d) 0

55) If $A = \begin{bmatrix} i & 0 & 0 \\ 0 & i & 0 \\ 0 & 0 & i \end{bmatrix}$, $i = \sqrt{-1}$, then $A^n = I$ where I is unit matrix when $n =$ _____

(a) $4p + 1$ (b) $4p + 3$ (c) $4p$ (d) $4p + 2$

56) If $A = \begin{bmatrix} k & 3 \\ 3 & k \end{bmatrix}$ and $|A^3| = 343$, then find the value of k _____

(a) ± 1 (b) ± 2 (c) ± 3 (d) ± 4

57) If $A = \begin{bmatrix} 1 & 0 \\ 1 & 1 \end{bmatrix}$, then $A^n + (n-1)I =$ _____

(a) $2^{n-1}A$ (b) $-nA$ (c) nA (d) $(n+1)A$

58) If $\begin{bmatrix} x^2+x & x+1 & x-2 \\ 2x^2+3x-1 & 3x & 2x-3 \\ x^2+2x+3 & 2x-1 & 2x-1 \end{bmatrix} = 24x + B$ then $B =$ _____

(a) -12 (b) 12 (c) 24 (d) -8

59) $X \begin{bmatrix} \tan^2 x & -\sec^2 x & 1 \\ -\sec^2 x & \tan^2 x & 1 \\ -10 & 12 & -2 \end{bmatrix} =$ _____

(a) $12 \tan^2 x - 10 \sec^2 x$ (b) $12 \sec^2 x - 10 \tan^2 x + 2$ (c) 0
(d) $\tan^2 x \cdot \sec^2 x$

60) If $f(x) = \begin{vmatrix} \sec x & \cos x & \sec^2 x + \cos x \operatorname{cosec}^2 x \\ \cos^2 x & \cos^2 x & \operatorname{cosec}^2 x \\ 1 & \cos^2 x & \operatorname{cosec}^2 x \end{vmatrix}$

then $\int_0^{\frac{\pi}{2}} f(x) dx =$ _____

(a) $\frac{1}{3} - \frac{\pi}{3}$ (b) $\frac{1}{3} - \frac{\pi}{4}$ (c) $\frac{2}{3} + \frac{\pi}{3}$ (d) $\frac{4}{3} - \frac{\pi}{4}$

61) If $f(x) = \begin{vmatrix} x & e^{x^2} & \sec x \\ \sin x & 2 & \cos x \\ \operatorname{cosec} x & x^2 & 5 \end{vmatrix}$, then the value of $\int_0^{\frac{\pi}{2}} f(x) dx =$ _____

(a) 0 (b) $5e^\pi$ (c) $1 - \frac{\pi}{2}$ (d) 34

62) If $A = \begin{bmatrix} a & b & c \\ b & c & a \\ c & a & b \end{bmatrix}$, where $a, b, c \in \mathbb{R}^+$, $abc = 1$ and $|A| > 0$, $A^T A = I$, then $a^3 + b^3 + c^3 =$ _____

(a) 12 (b) 4 (c) -8 (d) 24

63) The value of the determinant $\begin{vmatrix} \cos^2\left(\frac{\pi}{2} + x\right) & \cos^2\left(\frac{3\pi}{2} + x\right) & \cos^2\left(\frac{5\pi}{2} + x\right) \\ \cos\left(\frac{\pi}{2} + x\right) & \cos\left(\frac{3\pi}{2} + x\right) & \cos\left(\frac{5\pi}{2} + x\right) \\ \cos\left(\frac{\pi}{2} - x\right) & \cos\left(\frac{3\pi}{2} - x\right) & \cos\left(\frac{5\pi}{2} - x\right) \end{vmatrix}$ is _____

(a) 0 (b) $\cos^2\left(3x - \frac{9\pi}{2}\right)$ (c) $\sin^2\left(\frac{3\pi}{2} + x\right)$ (d) $\cos^2\left(\frac{15\pi}{2} - x\right)$

64) If $A = \begin{bmatrix} 3a & b & c \\ b & 3c & a \\ c & a & 3b \end{bmatrix}$, $a, b, c \in \mathbb{R}$, $abc = 1$ and $AA^T = 64I$ and $|A| > 0$, then $|a^3 + b^3 + c^3|^3 =$ _____

(a) 343 (b) 729 (c) 256 (d) 512

65) Let P be a non-singular matrix and $1 + P + P^2 + \dots + P^n = O$, (O denotes the null matrix) then $P^{-1} =$ _____

(a) 0 (b) P (c) P^n (d) I

66) If the inverse of the matrix $\begin{bmatrix} 1 & 2 \\ 3 & -5 \end{bmatrix}$ is $\frac{1}{11} \begin{bmatrix} a & b \\ c & d \end{bmatrix}$ then the ascending order of a, b, c, d is _____

(a) a, b, c, d (b) d, b, c, a (c) c, a, b, d (d) b, d, c, d

67) If A and B are orthogonal, then $(AB)^T (AB)$ is _____

(a) A (b) B (c) I (d) A^T

68) The adjoint of 3×3 matrix P is $\begin{bmatrix} -1 & 2 & 2 \\ 1 & 1 & 2 \\ 2 & 2 & 1 \end{bmatrix}$, then the possible value (s) of the determinant p is / are _____

(a) 3 (b) -3 (c) ± 3 (d) $\pm \sqrt{3}$

69) If A is a 3×3 matrix such that $|3 \text{adj } A| = 3$ then $|A|$ is equal to _____

(a) $\frac{1}{3}$ (b) $-\frac{1}{3}$ (c) $\pm \frac{1}{3}$ (d) ± 3

(Match the following)

20 x 1 = 20

70) Trivial solution of $AX = 0$ (1) $B^{-1}A^{-1}$

71) Non - Trivial solution of $AX = 0$ (2) Unique solution

72) $\rho(A) = \rho[(A|0)] < n$ (3)

Consistent with one parameter family of solution

73) $\rho(A) = \rho[(A|0)] = n$ (4) $|A|^{n-2} \cdot A$

74) $\rho(A) = \rho[(A|B)] = 3 = \text{number of unknowns}$ (5) $|A|^{n-1}$

75) $\rho(A) = \rho[(A|B)] = 2 < \text{number of unknowns}$ (6) $(\text{adj } B)(\text{adj } A)$

76) $\rho(A) = \rho[(A|B)] = 1 < \text{number of unknowns}$ (7) $|A| \cdot I_n$

77) $\rho(A) \neq \rho[(A|B)]$ (8) Non - trivial solution

78) $[\text{adj } A]$ (9) $\lambda^{n-1} \text{adj}(A)$

79) $(\text{adj } A)^T$ (10) Trivial solution

80) $\text{adj}(\text{adj } A)$ (11) $\text{adj}(A^{-1})$

81) $|\text{adj}(\text{adj } A)|$ (12) $|A| \neq 0$

82) $(\text{adj } A)^{-1}$ (13)

Consistent with two parameter family of solution

83) $(\lambda A)^{-1}$ (14) $\text{adj}(A^T)$

84) $\text{adj}(AB)$ (15) $|A|^{n-2}A$

85) $(A^T)^{-1}$ (16) $\frac{1}{\lambda}A^{-1}$

86) $A(\text{adj } A)$ (17) $(A^{-1})^T$

87) $(AB)^{-1}$ (18) In consistent and has no solution

88) $(A^{-1})^{-1}$ (19) A

89) $\text{adj}(\lambda A)$ (20) $|A| = 0$

(Odd one out)

90) The rank of any 3×4 matrix is

- (1) May be 1
- (2) May be 2
- (3) May be 3
- (4) Maybe 4

91) If A is symmetric then

- (1) $A^T = A$
- (2) $\text{adj } A$ is symmetric
- (3) $\text{adj}(A^T) = (\text{adj } A)^T$
- (4) A is orthogonal

92) If A is a non-singular matrix of odd order then

- 1) Order of A is $2m + 1$
- (2) Order of A is $2m + 2$
- (3) $|\text{adj } A|$ is positive
- (4) $|A| \neq 0$

JOIN 10TH & 12TH PAID WHATSAPP GROUP

DEC 1 2025 TO TILL FINAL 2026 MARCH EXAM
UPLOAD DAILY ONE SUBJECT FULL TEST PAPER

ONE TIME FEES RS.750 ONLY

ALL TEST PAPERS WITH ANSWERS UPLOAD IN YOUR WHATSAPP

FREE USERS CHECK ALL SAME PAPERS UPLOAD IN MY
WEBSITES. YOU CAN DOWNLOAD AND PRACTICE

RAVI TEST PAPERS & NOTES
WHATSAPP - 8056206308

CHECK FREE TEST PAPERS www.ravitestpapers.com & www.ravitestpapers.in

5 x 2 = 10

93) If A is a orthogonal matrix, then

- (1) $AA^T = A^T A = I$
- (2) A is non-singular
- (3) $|A| = 0$
- (4) $A^{-1} = A^T$

94) A matrix which is obtained from an identity matrix by applying only one elementary transformation is

- (1) Identity matrix
- (2) Elementary matrix
- (3) Square matrix
- (4) Equivalent to identity matrix

5 x 2 = 10

(Find the wrong statement)

95) In an echelon form which of the following is incorrect?

- (1) Every row of a which has all its entries 0 occurs below every row which has a non- zero entry.
- (2) The first non-zero entry in each non-zero row is 1
- (3) The number of zeros before the first non- zero element in a row is less than the number of such zeros in the next row
- (4) Two row can have same number of zeros before the first non-zero entry

96) Which of the following elementary transformation is not correct?

- (1) $R_i \rightarrow R_i + 2R_j$
- (2) $C_i \rightarrow C_i - C_j$
- (3) $R_i \rightarrow 7R_i + \frac{5}{3}R_j$
- (4) $C_i \rightarrow C_i - R_j$

97) If A is an invertible matrix, then which of the following is not true.

- (1) $(A^2)^{-1} = (A^{-1})^2$
- (2) $|A^{-1}| = |A|^{-1}$
- (3) $(A^T)^{-1} = (A^{-1})^T$
- (4) $A \neq 0$

98) The matrix $\begin{bmatrix} 5 & 10 & 3 \\ -2 & -4 & 6 \\ -1 & -2 & x \end{bmatrix}$ is a singular matrix if the value of x is

- (1) 3
- (2) non-existent
- (3) All values of x
- (4) Any value of x

99) The number of solutions of the system of equations $2x + y - z = 7$, $x - 3y + 2z = 1$, $x + 3y - 3z = 5$ is

- (1) 0
- (2) 3
- (3) No-solution
- (4) Inconsistent

(Multiple Choice Question)

69 x 1 = 69